

Uniform Euler approximation of solutions of fractional-order delayed cellular neural network on bounded intervals

Swati Tyagi¹, Syed Abbas^{1*}, Manuel Pinto² and Daniel Sepúlveda³

¹School of Basic Sciences, Indian Institute of Technology, Mandi, H.P., 175001, India.

²Departamento de Matemáticas, Facultad de Ciencias, Universidad de Chile, Casilla 653, Santiago, Chile.

³Escuela de Matemática y Estadística, Universidad Central de Chile, Toesca 1783, Santiago, Chile.

E-mail: sabbas.iitk@gmail.com*

Abstract

In this paper, we study convergence characteristics of a class of continuous time fractional-order cellular neural network containing delay. Using the method of Lyapunov and Mittag-Leffler functions, we derive sufficient condition for global Mittag-Leffler stability, which further implies global asymptotic stability of the system equilibrium. Based on the theory of fractional calculus and the generalized Gronwall inequality, we approximate the solution of the corresponding neural network model using discretization method by piecewise constant argument and obtain some easily verifiable conditions, which ensures that the discrete-time analogues preserve the convergence dynamics of the continuous-time networks. In the end, we give appropriate examples to validate the proposed results, illustrating advantages of the discrete-time analogue over continuous-time neural network for numerical simulation.

2010 Mathematics Subject Classification. **26A33.** 92B20; 34K07; 34A45

Keywords. Fractional-order; Neural network; Mittag-Leffler stability; Approximate solution; Error analysis; Piecewise constant argument of generalized type.

1 Introduction

The Fractional-calculus as a generalization of integer order differentiation and integral to non-integer order was first introduced 300 years ago, though their applications are rather recent. It has wide range of applications in control theory [1], viscoelasticity [2], signal processing [3] and social sciences [4].

It has been recognized that the integral derivative of a function is related only to its nearby points, while fractional derivative is related to all of the function history. It can be interpreted in other way that the future state of the system depends on its present state as well as upon its history values starting from the initial point. Thus fractional-order model possesses memory, and due to this, it becomes precise to describe the state of the neurons. Fractional derivative has ability to model memory and hereditary properties, since the fractional-derivative is non-local. Thus the models containing fractional derivative are very crucial. Recently, it has been found that the fractional-order calculus could be used in the study of Hopfield neural networks. Various results on fractional-order neural networks have been obtained as a result of development of fractional calculus [5–8]. The discrete time fractional-order artificial neural networks were presented in [9, 10].

*Corresponding author

Due to finite switching speeds of the amplifiers, time-delay is inevitable, which can also result in oscillations or instability of dynamic system. Thus, it becomes an interesting topic to study the stability of fractional-order neural networks containing delays. Also in some literatures, chaos and chaotic synchronization has been discussed [11–13].

In [14], Zhang et al. have investigated the Mittag-Leffler stability for a fractional-order neural network. In this paper, we study the Mittag-Leffler stability for a fractional-order neural network by introducing time delay term. Moreover, we derive sufficient approximating results for the solutions of fractional-order delayed neural network via solutions of corresponding fractional-order delayed neural network with piecewise constant arguments of generalized type. Having obtained sufficient conditions for the convergence dynamics of the continuous-time network, we propose an innovative method for obtaining discrete-time analogues from the continuous-time network. In particular, we propose a novel method to obtain a discrete-time fractional-order dynamical system whose dynamics is inherited from the continuous-time dynamic system. The discretization method proposed in this paper is different from the other discretization methods (such as predictor-corrector and Euler method), in the sense that our method is an approximation via piecewise constant arguments of generalized type, which has not been obtained in the related field of neural networks so far. Thus the results obtained are new and more general than the previous results derived in literature.

In this paper, we consider the following delayed fractional-order neural network model described by the following differential equation:

$$\begin{aligned} D_t^\alpha x_i(t) &= -a_i x_i(t) + \sum_{j=1}^m b_{ij} f(x_j(t)) + \sum_{j=1}^m c_{ij} g(x_j(t-\tau)) + I_i \quad t \in [0, T] \\ x_i(t) &= \varphi_i(t), \quad t \in [-\tau, 0], \quad T < \infty, \quad i = 1, 2, \dots, m, \end{aligned} \quad (1.1)$$

where m represents the number of units in the neural network; D_t^α denotes the Liouville-Caputa fractional derivative of order α , $0 < \alpha \leq 1$; $x(t) = (x_1(t), x_2(t), \dots, x_m(t))^T \in \mathbb{R}^m$ corresponds to the state vector at time t ; $f(\cdot)$ and $g(\cdot)$ denote the activation functions; I_i are constants denote the external bias on the i^{th} neuron; the positive constants $a_i > 0$ denote the rates with which the i^{th} neuron will reset its potential to the resting state in isolation when disconnected from the networks and external inputs; the constants b_{ij} and c_{ij} denote the connections of j^{th} neuron to i^{th} neuron at time t and $t - \tau$; τ is the transmission delay respectively; $a_i = \text{diag}(a_1, a_2, \dots, a_m)$, b_{ij} , c_{ij} are the connection weight matrices; and $I_i = (I_1, I_2, \dots, I_m)^T$ are external inputs. The initial conditions are $\varphi_i(t) \in C([-\tau, 0], \mathbb{R}^m)$, where $C([-\tau, 0], \mathbb{R}^m)$ denotes the Banach space of continuous m -real vector functions defined on $[-\tau, 0]$. $[p]$ denotes the integer part of p .

The structure of this paper is as follows. In Section 2, we give some preliminaries regarding fractional order differential systems. In Section 3, we establish the Mittag-Leffler stability of system (1.1) by constructing suitable Lyapunov function. In Section 4, using discretization scheme, we show that that the solution $z_i^h(t)$ of (4.2) approximates the solution $x_i(t)$ of (1.1) uniformly on any compact interval $[0, T]$, $T < \infty$. Later, we prove that the error function converges to the actual solution for $h \rightarrow 0$ uniformly. Moreover, in Section 5, we present some numerical results to illustrate the effectiveness of the derived results followed with conclusion in Section 6.

2 Preliminaries on fractional - order differential systems

For the convenience of reader, we recall some definitions and lemmas of fractional calculus in this section (for instance, see [15–17]).

Definition 2.1. [20] The Riemann-Liouville fractional integral with fractional-order $\alpha > 0$ of function $f(t)$ is defined as

$${}_{t_0}I_t^\alpha f(t) = \frac{1}{\Gamma(\alpha)} \int_{t_0}^t (t-s)^{\alpha-1} f(s) ds, \tag{2.1}$$

where $\Gamma(\cdot)$ is the Gamma function and is given by $\Gamma(z) = \int_0^\infty t^{z-1} e^{-t} dt$.

Definition 2.2. [19] The Liouville-Caputa derivative of fractional order α of a function $f(t)$ is defined as

$${}_C D_{t_0,t}^\alpha f(t) = \frac{1}{\Gamma(n-\alpha)} \int_{t_0}^t (t-s)^{n-\alpha-1} f^{(n)}(s) ds \tag{2.2}$$

where $n-1 < \alpha < n \in \mathbb{Z}^+$.

Remark 2.3. For sake of simplicity, we denote

$${}_C D_{t_0,t}^\alpha f(t) = D_t^\alpha f(t). \tag{2.3}$$

Remark 2.4. When $\alpha \rightarrow 1$, the fractional order derivative $D_t^\alpha f(t)$ converges to the integer order derivative $f'(t)$.

Definition 2.5. [19] The two-parameter Mittag-Leffler function is defined as:

$$E_{\alpha,\beta}(z) = \sum_{k=0}^\infty \frac{z^k}{\Gamma(k\alpha + \beta)}, \tag{2.4}$$

where $\alpha > 0, \beta > 0$ and $z \in \mathbb{C}$, where \mathbb{C} denotes the complex plane. When $\beta = 1$, we have

$$E_{\alpha,1}(z) = E_\alpha(z) = \sum_{k=0}^\infty \frac{z^k}{\Gamma(k\alpha + 1)} \tag{2.5}$$

Further, for $\alpha = 1, \beta = 1$, we have $E_{1,1}(z) = e^z$.

Lemma 2.6. [19] The following properties exist for the Mittag-Leffler function.

1. There exists constants $N_1, N_2 \geq 1$ such that for any $0 < \alpha < 1$,

$$\begin{aligned} \|E_{\alpha,1}(At^\alpha)\| &\leq N_1 \|e^{At}\|, \\ \|E_{\alpha,\alpha}(At^\alpha)\| &\leq N_2 \|e^{At}\|, \end{aligned} \tag{2.6}$$

where A denotes matrix, $\|\cdot\|$ denotes any vector or induced matrix norm.

2. If $\alpha \geq 1$, then for $\beta = 1, 2, \alpha$

$$\|E_{\alpha,\beta}(At^\alpha)\| \leq \|e^{At^\alpha}\|, \quad t \geq 0. \quad (2.7)$$

Consider an n -dimensional fractional-order system given by

$${}_t D_t^\alpha = f(t, x(t)), \quad x(t_0) = x_{t_0}. \quad (2.8)$$

Definition 2.7. (Generalized Mittag-Leffler stability [14]) The equilibrium point $x^* = (x_1^*, x_2^*, \dots, x_m^*)^T$ of system (2.8) is said to be globally Mittag-Leffler stable, if

$$\|x(t) - x^*\| \leq [m(x_{t_0} - x^*)E_\alpha(-\gamma(t - t_0)^\alpha)]^b,$$

where $\gamma > 0$, $b > 0$, $m(0) = 0$, $m(x) \geq 0$ and $m(x)$ satisfies locally Lipschitz condition on $x \in \mathbb{R}^m$ with Lipschitz constant m_0 .

Remark 2.8. The global Mittag-Leffler stability implies global asymptotic stability.

Theorem 2.9. (The Second Method of Lyapunov [14]) For $t_0 = 0$, the fractional-order system (2.8) is Mittag-Leffler stable at the equilibrium point $x^* = 0$, if there exists a continuously differentiable function $V(t, x(t))$ satisfying

$$\alpha_1 \|x\|^a \leq V(t, x(t)) \leq \alpha_2 \|x\|^{ab}, \quad (2.9)$$

$$D_t^\beta V(t, x(t)) \leq -\alpha_3 \|x\|^{ab}, \quad (2.10)$$

where $V(t, x(t)) : [0, \infty) \times D \rightarrow \mathbb{R}$ satisfies locally Lipschitz condition on x ; $D \subset \mathbb{R}^m$ is a domain containing origin; $t \geq 0$, $\beta \in (0, 1)$, $\alpha_1, \alpha_2, \alpha_3, a$ and b are arbitrary positive constants. If the assumptions hold globally on \mathbb{R}^m , then $x^* = 0$ is globally Mittag-Leffler stable.

Theorem 2.10. (The Extended Second Method of Lyapunov [14]) For $t_0 = 0$, the fractional-order system (2.8) is Mittag-Leffler stable at the equilibrium point $x^* = 0$, if there exists a continuous function $V(t, x(t))$ satisfying

$$\alpha_1 \|x(t)\|^a \leq V(t, x(t)) \leq \alpha_2 \|x(t)\|^{ab}, \quad (2.11)$$

$$D_t^\beta V(t^+, x(t^+)) \leq -\alpha_3 \|x\|^{ab}, \quad (\text{holding almost everywhere}) \quad (2.12)$$

where $V(t, x(t)) : [0, \infty) \times D \rightarrow \mathbb{R}$ satisfies locally Lipschitz condition on x ; $\dot{V}(t, x(t))$ is piecewise continuous, and $\lim_{s \rightarrow t^+} \dot{V}(s, x(s))$ exists for any $t \in [0, \infty)$; $D \subset \mathbb{R}^m$ is a domain containing origin and $V(t^+, x(t^+)) = \lim_{s \rightarrow t^+} V(s, x(s))$; $t \geq 0$, $\beta \in (0, 1)$, $\alpha_1, \alpha_2, \alpha_3, a$ and b are arbitrary positive constants. If the assumptions hold globally on \mathbb{R}^m , then $x^* = 0$ is globally Mittag-Leffler stable.

Theorem 2.11. [14] If $h(t) \in C^1([0, +\infty), \mathbb{R})$, then the following inequality holds almost everywhere.

$$D_t^\alpha |h(t^+)| \leq \text{sgn}(h(t)) D_t^\alpha h(t), \quad 0 < \alpha < 1, \quad (2.13)$$

where $h(t^+) = \lim_{s \rightarrow t^+} h(s)$, where sgn denotes the signum function.

Lemma 2.12. [20] Let m be a positive integer such that $m - 1 < \alpha < m$; if $x(t) \in C^{m-1}[a, b]$, then

$$I_t^\alpha D_t^\alpha x(t) = x(t) - \sum_{k=0}^{m-1} \frac{x^{(k)}(a)}{k!} (t-a)^k. \tag{2.14}$$

In particular, if $0 < \alpha \leq 1$ and $x(t) \in C[a, b]$, then we have the Fundamental Theorem of Calculus

$$I_t^\alpha D_t^\alpha x(t) = x(t) - x(a). \tag{2.15}$$

Lemma 2.13. [33] Suppose that $\alpha > 0$, $x(t)$ be a non-negative function locally integrable on $0 \leq t < T$ (for some $T \leq +\infty$) and v be a non-negative, non-decreasing continuous function defined on $0 \leq t < T$, $v(t) \leq M$ (real constant). Furthermore, if $u(t)$ is non-negative and locally integrable on $0 \leq t < T$ such that

$$x(t) \leq u(t) + v(t) \int_0^t (t-s)^{\alpha-1} x(s) ds,$$

on $0 \leq t < T$. Then

$$x(t) \leq u(t) + \int_0^t \left[\sum_{m=1}^{\infty} \frac{(v(t)\Gamma(\alpha))^m}{\Gamma(m\alpha)} (t-s)^{m\alpha-1} u(s) \right] ds, \quad t \in [0, T]. \tag{2.16}$$

Henry [18] obtained (2.16) for u and v constant. For the use of (2.16), see [19].

Remark 2.14. If in Lemma 2.13, $u(t)$ is a non-decreasing function on $0 \leq t < T$, then

$$x(t) \leq u(t) E_{\alpha,1}(v(t)\Gamma(\alpha)t^\alpha), \quad t \in [0, T]. \tag{2.17}$$

3 Global Mittag-Leffler stability

In this section, we obtain sufficient condition for global Mittag-Leffler stability of the system (1.1), which further implies the global asymptotic stability of the equilibrium point.

We introduce the following conditions on our model system (1.1).

- A1. The activation functions $f(\cdot)$ and $g(\cdot)$ are continuous and Lipschitz, i.e. for some $x, y \in \mathbb{R}$, there exist positive constants L_f and L_g such that

$$\begin{aligned} |f(x) - f(y)| &\leq L_f |x - y|, \\ |g(x) - g(y)| &\leq L_g |x - y|. \end{aligned}$$

- A2. There exists positive constant $c_i (i = 1, 2, \dots, m)$ such that

$$c_i = \frac{1}{a_i} \left(\sum_{j=1}^m |b_{ji}| L_f + \sum_{j=1}^m |c_{ji}| L_g \right) < 1. \tag{3.1}$$

A3. The system parameters a_i, b_{ij}, c_{ij} ($i, j = 1, 2, \dots, m$) satisfy

$$\min_{1 \leq i \leq m} \left(a_i - L_f \sum_{j=1}^m |b_{ji}| \right) > \max_{1 \leq i \leq m} \left(L_g \sum_{j=1}^m |c_{ji}| \right) > 0. \tag{3.2}$$

A4. The time delay term τ is a positive real number.

Remark 3.1. The problems of existence and uniqueness of equilibrium solutions of fractional order delayed neural networks have been studied by several authors in literature. For more details, refer to [20–22]. For convenience of reader, we give the proof of existence and uniqueness of equilibrium point of system (1.1) as following theorem.

Theorem 3.2. If the Assumptions (A1)-(A2) hold, then there exists a unique equilibrium point of system (1.1).

Proof. Define $\|x\| = \|x\|_1$, that is $\|x\| = \sum_{i=1}^m |x_i|$ for any $x = (x_1, x_2, \dots, x_m)^T \in \mathbb{R}^m$. Consider the mapping

$F(y) = (F_1(y_1), F_2(y_2), \dots, F_m(y_m))^T$, where $y = (y_1, y_2, \dots, y_m)^T \in \mathbb{R}^m$ and

$$F_i(y_i) = \sum_{j=1}^m b_{ij} f\left(\frac{y_j}{a_j}\right) + \sum_{j=1}^m c_{ij} g\left(\frac{y_j}{a_j}\right) + I_i, \quad (i = 1, 2, \dots, m).$$

We will show that F is a contraction mapping on \mathbb{R}^m . From Assumption (A1), for any two vectors $y, z \in \mathbb{R}^m$, we have

$$\begin{aligned} |F_i(y) - F_i(z)| &= \left| \sum_{j=1}^m b_{ij} \left(f\left(\frac{y_j}{a_j}\right) - f\left(\frac{z_j}{a_j}\right) \right) \right| + \left| \sum_{j=1}^m c_{ij} \left(g\left(\frac{y_j}{a_j}\right) - g\left(\frac{z_j}{a_j}\right) \right) \right| \\ &\leq \left(\sum_{j=1}^m \frac{|b_{ij}|L_f}{a_j} + \sum_{j=1}^m \frac{|c_{ij}|L_g}{a_j} \right) |y_j - z_j| \end{aligned}$$

From Assumption (A2), we obtain

$$\begin{aligned} \|F(y) - F(z)\| &= \sum_{i=1}^m |F_i(y) - F_i(z)| \\ &\leq \sum_{i=1}^m \sum_{j=1}^m \frac{1}{a_j} (|b_{ij}|L_f + |c_{ij}|L_g) |y_j - z_j| \\ &= \sum_{i=1}^m \left(\sum_{j=1}^m \frac{1}{a_i} (|b_{ji}|L_f + |c_{ji}|L_g) \right) |y_i - z_i| \\ &< c \|y - z\|, \end{aligned}$$

where $c = \max\{c_1, c_2, \dots, c_m\}$. Thus we have $\|F(y) - F(z)\| < c\|y - z\|$. Hence the mapping $F : \mathbb{R}^m \rightarrow \mathbb{R}^m$ is a strict contraction mapping on \mathbb{R}^m with Lipschitz constant c . Thus there exists a unique fixed point $y^* \in \mathbb{R}^m$ such that $F(y^*) = y^*$, that is

$$y_i^* = \sum_{j=1}^m b_{ij} f\left(\frac{y_j^*}{a_j}\right) + \sum_{j=1}^m c_{ij} g\left(\frac{y_j^*}{a_j}\right) + I_i, \quad (i = 1, 2, \dots, m).$$

Denote $x_i^* = \frac{y_i^*}{a_i}$ for $i = 1, 2, \dots, m$. Then we have

$$-a_i x_i^* + \sum_{j=1}^m b_{ij} f(x_j^*) + \sum_{j=1}^m c_{ij} g(x_j^*) + I_i = 0, \quad (i = 1, 2, \dots, m).$$

Thus $x^* = (x_1^*, x_2^*, \dots, x_m^*)^T$ is the unique equilibrium point of system (1.1). Q.E.D.

Now we prove the Mittag-Leffler stability of the unique equilibrium point of system (1.1).

Let $x^* = (x_1^*, x_2^*, \dots, x_m^*)^T$ be the unique equilibrium point of (1.1). Then for $e \in \mathbb{R}^m$, $e = (e_1, e_2, \dots, e_m)^T$, it can be derived from (1.1) that the error $e_i(t) = x_i(t) - x_i^*$ satisfies

$$\begin{aligned} D_t^\alpha e_i(t) &= -a_i e_i(t) + \sum_{j=1}^m b_{ij} (f(x_j^* + e_j(t)) - f(x_j^*)) \\ &\quad + \sum_{j=1}^m c_{ij} (g(x_j^* + e_j(t - \tau)) - g(x_j^*)) \\ e_i(t) &= \varphi_i(t) - x_i^*, \quad t \in [-\tau, 0] \quad i = 1, 2, \dots, m. \end{aligned} \quad (3.3)$$

Theorem 3.3. If Assumptions (A1)–(A4) hold, then the equilibrium point $x^* = (x_1^*, x_2^*, \dots, x_m^*)^T$ of system (1.1) is globally Mittag-Leffler stable.

Proof. We construct a Lyapunov function as

$$V(t, e(t)) = \|e(t)\| = \sum_{i=1}^m |e_i(t)|. \quad (3.4)$$

It can be observed that the Lyapunov function satisfies equation (2.9) of Theorem 2.10. From

Assumptions (A1), (A3) and Theorem 2.11, the following inequality holds almost everywhere;

$$\begin{aligned}
 D_t^\alpha V(t^+, e(t^+)) &= \sum_{i=1}^m D_t^\alpha |e_i(t^+)| \leq \sum_{i=1}^m \operatorname{sgn}(e_i(t)) D_t^\alpha e_i(t) \\
 &\leq \sum_{i=1}^m \left[-a_i |e_i(t)| + \sum_{j=1}^m L_f |b_{ij}| |e_j(t)| + \sum_{j=1}^m L_g |c_{ij}| |e_j(t - \tau)| \right] \\
 &= -\sum_{i=1}^m \left[a_i - \sum_{j=1}^m L_f |b_{ji}| \right] |e_i(t)| + \sum_{j=1}^m \sum_{i=1}^m L_g |c_{ij}| |e_j(t - \tau)| \\
 &\leq -k_1 V(t, e(t)) + k_2 \sup_{t-\tau \leq s \leq t} V(s, e(s)),
 \end{aligned}$$

where

$$k_1 = \min_{1 \leq i \leq m} \left(a_i - L_f \sum_{j=1}^m |b_{ji}| \right) > 0, \text{ and } k_2 = \max_{1 \leq i \leq m} \left(L_g \sum_{j=1}^m |c_{ji}| \right) > 0. \tag{3.5}$$

Thus from above estimate, for any solution $e(t)$ of (3.3), that satisfies the Razumikhin condition

$$V(s, e(s)) \leq V(t, e(t)), \quad t - \tau \leq s \leq t,$$

we have

$$D_t^\alpha V(t^+, e(t^+)) \leq -(k_1 - k_2)V(t, e(t)). \tag{3.6}$$

Form Assumption (A3), there exists a real number $\gamma > 0$, such that

$$k_1 - k_2 \geq \gamma.$$

Thus it follows that

$$D_t^\alpha V(t^+, e(t^+)) \leq -\gamma V(t, e(t)). \tag{3.7}$$

It can be observed that the condition (2.12) of Theorem 2.10 is satisfied. Thus, we have

$$\|e(t)\| = V(t, e(t)) \leq V(0, e(0))E_\alpha(-\gamma t^\alpha). \tag{3.8}$$

Hence,

$$\begin{aligned}
 \|x(t) - x^*\| &= \sum_{i=1}^m |x_i(t) - x_i^*| \leq E_\alpha(-\gamma t^\alpha) \sum_{i=1}^m |x_i(0) - x_i^*| \\
 &= \|\varphi_0 - x^*\| E_\alpha(-\gamma t^\alpha) \\
 &\leq \|\varphi_0 - x^*\|_\tau E_\alpha(-\gamma t^\alpha).
 \end{aligned}$$

Let $m = \|\varphi_0 - x^*\|_\tau$, where $m \geq 0$ and $m = 0$ holds only if $\varphi_i(0) = x_i^*$, ($i = 1, 2, \dots, m$). Then $\|x(t) - x^*\| \leq mE_\alpha(-\gamma t^\alpha)$. Thus system (1.1) is globally Mittag-Leffler stable. Q.E.D.

Remark 3.4. Since the equilibrium point x^* of system (1.1) is globally Mittag-Leffler stable. Thus from Remark 2.8, it is globally asymptotically stable.

4 Approximation Results

4.1 Formulation of discrete-time analogue: Euler's polygonal lines of order α

Recently, many researchers have applied the discretization method to discretize various cellular neural networks [23–29]. But discretizing the fractional-order differential equations is of main interest. In [30], Riccati's fractional-order differential equation is discretized. It has been observed that as the fractional-order parameter $\alpha \rightarrow 1$, Euler's discretization method is obtained. In [31], logistic fractional-order differential equation is discretized using same method. In [32], discretization method is applied to Chua's system, which is one of the autonomous differential equations capable of generating chaotic behavior. Fixed points and their asymptotic stability are also investigated. Chaotic attractor, bifurcation and chaos for different values of the fractional-order parameter have been discussed. In the next result, we propose a numerical discretization scheme using piecewise constant arguments of generalized type, which has not been derived till now to the best of author's knowledge.

Let $0 < \alpha \leq 1$, and consider the following fractional-order delayed neural network.

$$\begin{aligned} D_t^\alpha x_i(t) &= -a_i x_i(t) + \sum_{j=1}^m b_{ij} f(x_j(t)) + \sum_{j=1}^m c_{ij} g(x_j(t - \tau)) + I_i \quad t \in [0, T] \\ x_i(t) &= \varphi_i(t), \quad t \in [-\tau, 0], \quad T < \infty, \quad i = 1, 2, \dots, m. \end{aligned} \quad (4.1)$$

Let $\gamma_h(t) = h \left\lceil \frac{t}{h} \right\rceil$, where $\lceil \cdot \rceil$ is the usual greatest integer function and h is a positive number in the interval $(0, \tau]$. In fact we can consider $h = \frac{\tau}{\nu}$ with $\nu \geq 1$ is an integer. It is clear that $\gamma_h(t) = hn$ for $hn \leq t \leq h(n+1)$, and n is a positive integer, also we have $\gamma_h(\tau) = \nu h$.

The corresponding differential equation with piecewise constant arguments

$\gamma_h(t) = h \left\lceil \frac{t}{h} \right\rceil$ is given by

$$\begin{aligned} D_t^\alpha z_i^h(t) &= -a_i z_i^h(\gamma_h(t)) + \sum_{j=1}^m b_{ij} f(z_j^h(\gamma_h(t))) \\ &+ \sum_{j=1}^m c_{ij} g(z_j^h(\gamma_h(t - \tau))) + I_i, \end{aligned} \quad (4.2)$$

with initial conditions

$$z_i^h(nh) = \varphi_i(nh), \quad n = -\nu, \dots, 0. \quad (4.3)$$

By the solution of above system (4.2)-(4.3), we mean a function $z^h = (z_i^h)$, $i = 1, 2, \dots, m$ defined on $\{-jh : j = 0, 1, \dots, \text{such that } -\tau \leq -jh \leq 0\}$ by (4.3), which satisfy the following properties on \mathbb{R}^+ ;

1. The functions z_i^h are continuous for each $i = 1, 2, \dots, m$ on \mathbb{R}^+ .

2. The corresponding derivatives of $z_i^h(t)$ exist at each point $t \in \mathbb{R}^+$ with the possible exception of the points $jh (j = 0, 1, \dots)$ where finite one-sided derivatives exists, for each $i = 1, 2, \dots, m$.
3. The function z_i^h satisfies (4.2) on each interval $I_{(j,h)} = [jh, (j+1)h)$ for $j = 0, 1, \dots$.

On $I_{(n,h)}$ we can rewrite equation (4.2)-(4.3) as

$$D_t^\alpha z_i^h(t) = -a_i z_i^h(nh) + \sum_{j=1}^m b_{ij} f(z_j^h(nh)) + \sum_{j=1}^m c_{ij} g(z_j^h(h(n-\nu))) + I_i. \quad (4.4)$$

As γ_h has been applied also in the linear part, the discretization method obtained is the α -fractional polygonal of Euler [30]. Our discretization method proceed as follows:

1. Let $t \in [0, h)$, then $\frac{t}{h} \in [0, 1)$. Thus we get, for $t \in [0, h)$

$$D_t^\alpha z_i^h(t) = -a_i z_i^h(0) + \sum_{j=1}^m b_{ij} f(z_j^h(0)) + \sum_{j=1}^m c_{ij} g(z_j^h(h(0-\nu))) + I_i,$$

and by the Fundamental Theorem of Fractional Calculus (2.15), the solution of (4.4) is given by

$$\begin{aligned} z_i^h(t) &= z_i^h(0) + \frac{1}{\Gamma(\alpha)} \int_0^t (t-s)^{\alpha-1} \left(-a_i z_i^h(0) + \sum_{j=1}^m b_{ij} f(z_j^h(0)) \right. \\ &\quad \left. + \sum_{j=1}^m c_{ij} g(z_j^h(h(0-\nu))) + I_i \right) ds, \end{aligned}$$

or

$$\begin{aligned} z_i^h(t) &= z_i^h(0) + \frac{t^\alpha}{\Gamma(\alpha+1)} \left(-a_i z_i^h(0) + \sum_{j=1}^m b_{ij} f(z_j^h(0)) \right. \\ &\quad \left. + \sum_{j=1}^m c_{ij} g(z_j^h(h(0-\nu))) + I_i \right), \end{aligned}$$

and by the continuity of $z_i^h(t)$ at $t = h$, we obtain

$$\begin{aligned} z_i^h(h) &= \left(1 - a_i \frac{h^\alpha}{\Gamma(\alpha+1)} \right) z_i^h(0) \\ &\quad + \frac{h^\alpha}{\Gamma(\alpha+1)} \left(\sum_{j=1}^m b_{ij} f(z_j^h(0)) + \sum_{j=1}^m c_{ij} g(z_j^h(h(0-\nu))) + I_i \right). \end{aligned} \quad (4.5)$$

2. Let $t \in [h, 2h)$, then $\frac{t}{h} \in [1, 2)$. Then we have

$$D_t^\alpha z_i^h(t) = -a_i z_i^h(t) + \sum_{j=1}^m b_{ij} f(z_j^h(t)) + \sum_{j=1}^m c_{ij} g(z_j^h(h(1-\nu))) + I_i.$$

The solution on $[h, 2h)$ of above is given as

$$\begin{aligned} z_i^h(t) &= z_i^h(h) + \frac{1}{\Gamma(\alpha)} \int_h^t (t-s)^{\alpha-1} \left(-a_i z_i^h(s) + \sum_{j=1}^m b_{ij} f(z_j^h(s)) \right. \\ &\quad \left. + \sum_{j=1}^m c_{ij} g(z_j^h(h(1-\nu))) + I_i \right) ds, \end{aligned}$$

or

$$\begin{aligned} z_i^h(t) &= z_i^h(h) - a_i \frac{(t-h)^\alpha}{\Gamma(\alpha+1)} z_i^h(h) \\ &\quad + \frac{(t-h)^\alpha}{\Gamma(\alpha+1)} \left(\sum_{j=1}^m b_{ij} f(z_j^h(h)) + \sum_{j=1}^m c_{ij} g(z_j^h(h(1-\nu))) + I_i \right), \end{aligned}$$

and by the continuity of $z_i^h(t)$ at $t = 2h$, we obtain

$$\begin{aligned} z_i^h(2h) &= \left(1 - a_i \frac{h^\alpha}{\Gamma(\alpha+1)} \right) z_i^h(h) \\ &\quad + \frac{h^\alpha}{\Gamma(\alpha+1)} \left(\sum_{j=1}^m b_{ij} f(z_j^h(h)) + \sum_{j=1}^m c_{ij} g(z_j^h(h(1-\nu))) + I_i \right). \end{aligned} \tag{4.6}$$

Using (4.5), we can rewrite (4.6) as

$$\begin{aligned} z_i^h(2h) &= \left(1 - a_i \frac{h^\alpha}{\Gamma(\alpha+1)} \right) \left(1 - a_i \frac{h^\alpha}{\Gamma(\alpha+1)} \right) z_i^h(0) \\ &\quad + \left(1 - a_i \frac{h^\alpha}{\Gamma(\alpha+1)} \right) \frac{h^\alpha}{\Gamma(\alpha+1)} \left(\sum_{j=1}^m b_{ij} f(z_j^h(0)) \right. \\ &\quad \left. + \sum_{j=1}^m c_{ij} g(z_j^h(h(1-\nu))) + I_i \right) \\ &\quad + \frac{h^\alpha}{\Gamma(\alpha+1)} \left(\sum_{j=1}^m b_{ij} f(z_j^h(h)) + \sum_{j=1}^m c_{ij} g(z_j^h(h(1-\nu))) + I_i \right). \end{aligned}$$

Repeating the process, it can be easily deduced that for $nh \leq t < (n+1)h$, the solution of IVP

(4.4) can be given as

$$\begin{aligned} z_i^h(t) &= z_i^h(nh) - a_i \frac{(t - nh)^\alpha}{\Gamma(\alpha + 1)} z_i^h(nh) \\ &+ \frac{(t - nh)^\alpha}{\Gamma(\alpha + 1)} \left(\sum_{j=1}^m b_{ij} f(z_j^h(nh)) + \sum_{j=1}^m c_{ij} g(z_j^h(h(n - \nu))) + I_i \right). \end{aligned} \quad (4.7)$$

Replacing $t = (n + 1)h$ in (4.7) and using continuity of z_i^h , we obtain

$$\begin{aligned} z_i^h((n + 1)h) &= \left(1 - a_i \frac{h^\alpha}{\Gamma(\alpha + 1)} \right) z_i^h(nh) \\ &+ \frac{h^\alpha}{\Gamma(\alpha + 1)} \left(\sum_{j=1}^m b_{ij} f(z_j^h(nh)) + \sum_{j=1}^m c_{ij} g(z_j^h(h(n - \nu))) + I_i \right). \end{aligned} \quad (4.8)$$

Define sequence $\mathbf{a}_i^h(n) = z_i^h(nh)$. This sequence satisfies the following non-linear system of delay difference equations

$$\begin{aligned} \mathbf{a}_i^h(n + 1) &= \left(1 - a_i \frac{h^\alpha}{\Gamma(\alpha + 1)} \right) \mathbf{a}_i^h(n) \\ &+ \frac{h^\alpha}{\Gamma(\alpha + 1)} \left(\sum_{j=1}^m b_{ij} f(\mathbf{a}_j^h(n)) + \sum_{j=1}^m c_{ij} g(\mathbf{a}_j^h(n - \nu)) + I_i \right). \end{aligned} \quad (4.9)$$

with initial condition

$$\mathbf{a}_i^h(n) = \varphi_i(nh) \quad n = -\nu, \dots, 0. \quad (4.10)$$

The initial value problem (4.9)-(4.10) is a discretization of the original problem (4.1).

From the above recurrence relation and initial conditions, we obtain

$$\mathbf{a}_i^h(0) = \varphi_i(0) \quad (4.11)$$

$$\begin{aligned} \mathbf{a}_i^h(1) &= \left(1 - a_i \frac{h^\alpha}{\Gamma(\alpha + 1)}\right) \mathbf{a}_i^h(0) \\ &\quad + \frac{h^\alpha}{\Gamma(\alpha + 1)} \left(\sum_{j=1}^m b_{ij} f(\mathbf{a}_j^h(0)) + \sum_{j=1}^m c_{ij} g(\mathbf{a}_j^h(0 - \nu)) + I_i \right) \\ &= \left(1 - a_i \frac{h^\alpha}{\Gamma(\alpha + 1)}\right) \varphi_i(0) \end{aligned} \quad (4.12)$$

$$\begin{aligned} &\quad + \frac{h^\alpha}{\Gamma(\alpha + 1)} \left(\sum_{j=1}^m b_{ij} f(\varphi_j(0)) + \sum_{j=1}^m c_{ij} g(\varphi_j(0 - \nu)) + I_i \right) \\ \mathbf{a}_i^h(2) &= \left(1 - a_i \frac{h^\alpha}{\Gamma(\alpha + 1)}\right) \mathbf{a}_i^h(1) \\ &\quad + \frac{h^\alpha}{\Gamma(\alpha + 1)} \left(\sum_{j=1}^m b_{ij} f(\mathbf{a}_j^h(1)) + \sum_{j=1}^m c_{ij} g(\mathbf{a}_j^h(1 - \nu)) + I_i \right) \\ &= \left(1 - a_i \frac{h^\alpha}{\Gamma(\alpha + 1)}\right) \left(1 - a_i \frac{h^\alpha}{\Gamma(\alpha + 1)}\right) \varphi_i(0) \\ &\quad + \left(1 - a_i \frac{h^\alpha}{\Gamma(\alpha + 1)}\right) \frac{h^\alpha}{\Gamma(\alpha + 1)} \left(\sum_{j=1}^m b_{ij} f(\varphi_j(0)) + \sum_{j=1}^m c_{ij} g(\varphi_j(0 - \nu)) + I_i \right) \\ &\quad + \frac{h^\alpha}{\Gamma(\alpha + 1)} \left(\sum_{j=1}^m b_{ij} f(\mathbf{a}_j^h(1)) + \sum_{j=1}^m c_{ij} g(\varphi_j(1 - \nu)) + I_i \right). \end{aligned} \quad (4.13)$$

Based on the above calculations, we can deduce that

$$\begin{aligned} \mathbf{a}_i^h(n+1) &= \left(1 - a_i \frac{h^\alpha}{\Gamma(\alpha + 1)}\right)^{(n+1)} \varphi_i(0) \\ &\quad + \sum_{k=0}^n \left(1 - a_i \frac{h^\alpha}{\Gamma(\alpha + 1)}\right)^{(n-k)} \frac{h^\alpha}{\Gamma(\alpha + 1)} \left[\sum_{j=1}^m b_{ij} f(\mathbf{a}_j^h(k)) \right. \\ &\quad \left. + \sum_{j=1}^m c_{ij} g(\mathbf{a}_j^h(k - \nu)) + I_i \right] \end{aligned} \quad (4.14)$$

From the initial conditions, the above sequence $\mathbf{a}_i^h(n)$ is well defined. Hence the solution of system

(4.3), for $t \in I_{(n,h)}$, can be written as

$$\begin{aligned}
 z_i^h(t) &= \mathbf{a}_i^h(n) + \frac{1}{\Gamma\alpha} \int_{nh}^t (t-s)^{\alpha-1} \left(-a_i \mathbf{a}_i^h(n) + \sum_{j=1}^m b_{ij} f(\mathbf{a}_j^h(n)) \right. \\
 &\quad \left. + \sum_{j=1}^m c_{ij} g(\mathbf{a}_j^h(n-\nu)) + I_i \right) ds
 \end{aligned} \tag{4.15}$$

are the α -fractional Euler method. We can write the above result as the following theorem.

Theorem 4.1. The system (4.2)-(4.3) has a unique solution, for $t \in I_{(n,h)}$, in the following form:

$$\begin{aligned}
 z_i^h(t) &= \left[1 - a_i \left(\frac{(t-nh)^\alpha}{\Gamma(\alpha+1)} \right) \right] \mathbf{a}_i^h(n) \\
 &\quad + \left(\frac{(t-nh)^\alpha}{\Gamma(\alpha+1)} \right) \left(\sum_{j=1}^m b_{ij} f(\mathbf{a}_j^h(n)) + \sum_{j=1}^m c_{ij} g(\mathbf{a}_j^h(n-\nu)) + I_i \right).
 \end{aligned} \tag{4.16}$$

for $t \geq 0$, $i = 1, 2, \dots, m$ and the sequence $\mathbf{a}_i^h(\cdot)$ satisfies the non linear difference equation (4.9) with the initial conditions given by (4.10).

Note 4.2. The approximation method with piecewise constant argument described above is valid only if the solutions of system of non-linear delayed differential equations (4.1), (4.3) and (4.9)-(4.10) are unique.

4.2 Approximation Results in Bounded Intervals: Error function

In this section, it is proved that system (4.1) can be approximated for $t \in [0, T)$ by the Euler schema.

To prove the result, we use the particular form of Lemma 2.12, i.e. for $0 < \alpha \leq 1$ and $x(t) \in C[a, b]$,

$$I_t^\alpha D_t^\alpha x(t) = x(t) - x(a). \tag{4.17}$$

Let there exists a positive constant r . Then for every $h \in (0, r)$ and $t \in I_{(n,h)} = [nh, (n+1)h), n \geq 0$, we define

$$\varepsilon_i^h(t) := x_i(t) - z_i^h(t). \tag{4.18}$$

We can rewrite (4.18) as

$$\begin{aligned}
 \varepsilon_i^h(t) &= x_i(t) - z_i^h(t) - \varphi_i(0) + \varphi_i(0) \\
 &= [x_i(t) - \varphi_i(0)] - [z_i^h(t) - \varphi_i(0)] \\
 &= I_t^\alpha D_t^\alpha x_i(t) - I_t^\alpha D_t^\alpha z_i^h(t).
 \end{aligned} \tag{4.19}$$

Since from Definition (2.1), we have

$$I_t^\alpha f(t) = \frac{1}{\Gamma\alpha} \int_0^t (t-s)^{\alpha-1} f(s) ds. \tag{4.20}$$

Using (4.20) in equation (4.19), we obtain

$$\begin{aligned}
 \varepsilon_i^h(t) &= \frac{1}{\Gamma\alpha} \int_0^t (t-s)^{\alpha-1} D_t^\alpha [x_i(s) - z_i^h(s)] ds \\
 &= \frac{-a_i}{\Gamma\alpha} \int_0^t (t-s)^{\alpha-1} [x_i(s) - z_i^h(\gamma_h(s))] ds \\
 &\quad + \frac{1}{\Gamma\alpha} \int_0^t (t-s)^{\alpha-1} \sum_{j=1}^m b_{ij} [f(x_j(s)) - f(z_j^h(\gamma_h(s)))] ds \\
 &\quad + \frac{1}{\Gamma\alpha} \int_0^t (t-s)^{\alpha-1} \sum_{j=1}^m c_{ij} [g(x_j(s-\tau)) - g(z_j^h(\gamma_h(s-\tau)))] ds.
 \end{aligned}
 \tag{4.21}$$

Note that

$$|x_i(s) - z_i^h(\gamma_h(s))| \leq |x_i(s) - x_i(\gamma_h(s))| + |x_i(\gamma_h(s)) - z_i^h(\gamma_h(s))|.$$

Using Lipschitz continuity of $f(\cdot)$ and $g(\cdot)$, we have

$$\begin{aligned}
 |\varepsilon_i^h(t)| &\leq \frac{a_i}{\Gamma\alpha} \int_0^t (t-s)^{\alpha-1} |\varepsilon_i^h(\gamma_h(s))| ds \\
 &\quad + \frac{a_i}{\Gamma\alpha} \int_0^t (t-s)^{\alpha-1} |x_i(s) - x_i(\gamma_h(s))| ds \\
 &\quad + \frac{L_f}{\Gamma\alpha} \int_0^t (t-s)^{\alpha-1} \sum_{j=1}^m |b_{ij}| |\varepsilon_j^h(\gamma_h(s))| ds \\
 &\quad + \frac{L_f}{\Gamma\alpha} \int_0^t (t-s)^{\alpha-1} \sum_{j=1}^m |b_{ij}| |x_i(s) - x_i(\gamma_h(s))| ds \\
 &\quad + L_g \frac{1}{\Gamma\alpha} \int_0^t (t-s)^{\alpha-1} \sum_{j=1}^m |c_{ij}| |\varepsilon_j^h(\gamma_h(s-\tau))| ds \\
 &\quad + L_g \frac{1}{\Gamma\alpha} \int_0^t (t-s)^{\alpha-1} \sum_{j=1}^m |c_{ij}| |x_j(s-\tau) - x_j(\gamma_h(s-\tau))| ds.
 \end{aligned}$$

Now we take the sum of the above inequalities from 1 to m , and obtain

$$\begin{aligned}
 & \sum_{i=1}^m |\varepsilon_i^h(t)| \\
 & \leq \sum_{i=1}^m \frac{a_i}{\Gamma\alpha} \int_0^t (t-s)^{\alpha-1} \left(\sum_{i=1}^m |\varepsilon_i^h(\gamma_h(s))| \right) ds \\
 & + \frac{L_f}{\Gamma\alpha} \int_0^t (t-s)^{\alpha-1} \sum_{1 \leq i,j \leq m} |b_{ij}| \left(\sum_{i=1}^m |\varepsilon_i^h(\gamma_h(s))| \right) ds \\
 & + \frac{L_g}{\Gamma\alpha} \int_0^t (t-s)^{\alpha-1} \sum_{1 \leq i,j \leq m} |c_{ij}| \left(\sum_{i=1}^m |\varepsilon_i^h(\gamma_h(s-\tau))| \right) ds \\
 & + \sum_{i=1}^m \frac{a_i}{\Gamma\alpha} \int_0^t (t-s)^{\alpha-1} \left(\sum_{i=1}^m |x_i(s) - x_i(\gamma_h(s))| \right) ds \\
 & + \frac{L_f}{\Gamma\alpha} \int_0^t (t-s)^{\alpha-1} \sum_{1 \leq i,j \leq m} |b_{ij}| \left(\sum_{i=1}^m |x_i(s) - x_i(\gamma_h(s))| \right) ds \\
 & + \frac{L_g}{\Gamma\alpha} \int_0^t (t-s)^{\alpha-1} \sum_{1 \leq i,j \leq m} |c_{ij}| \left(\sum_{i=1}^m |x_i(s-\tau) - x_i(\gamma_h(s-\tau))| \right) ds.
 \end{aligned} \tag{4.22}$$

Define $m(t) = \sup_{-\tau \leq s \leq t} \sum_{i=1}^m |\varepsilon_i^h(s)|$ and the *module of continuity* $w(x, h, T) = \max\{\sum_{i=1}^m |x_i(s) - x_i(\gamma_h(s))| : -\tau \leq s \leq T\}$. Therefore, we can write equation (4.22) for $t \geq 0$ as

$$m(t) \leq u(t) + \frac{\beta}{\Gamma\alpha} \int_0^t (t-s)^{\alpha-1} m(s) ds, \tag{4.23}$$

where

$$\beta := \sum_{i=1}^m a_i + \sum_{1 \leq i,j \leq m} (|b_{ij}|L_f + |c_{ij}|L_g), \tag{4.24}$$

and

$$u(t) := \frac{\beta t^\alpha}{\Gamma(\alpha + 1)} w(x, h, T).$$

Since $u(t)$ is non negative, non-decreasing and locally integrable function on $[0, T]$, β is a positive constant, then by Lemma 2.13, (2.17) holds, i.e.

$$m(t) \leq \frac{\beta t^\alpha}{\Gamma(\alpha + 1)} w(x, h, T) E_{\alpha,1}(\beta t^\alpha), \quad t \in [0, T].$$

Using (2.7) we obtain

$$m(t) \leq \frac{\beta t^\alpha}{\Gamma(\alpha + 1)} w(x, h, T) e^{\beta t}, \quad t \in [0, T].$$

From the hypothesis of theorem, it follows that on $[0, T]$, $\varepsilon_i^h(t) \rightarrow 0$ uniformly as $h \rightarrow 0$. We can summarize the above result as the following theorem.

Theorem 4.3. Assume that the assumptions (A1) and (A2) holds. Then for every $h \in (0, r)$, the solution x of the system (1.1) can be approximated using constant argument and the approximation satisfies:

$$|x_i(t) - z_i^h(t)| \leq \frac{\beta t^\alpha e^{\beta t}}{\Gamma(\alpha + 1)} w(x, h, T), \quad (4.25)$$

where β is given by (4.24) and $w(x, h, T) = \max\{\sum_{i=1}^m |x_i(s) - x_i(\gamma_h(s))| : -\tau \leq s \leq T\}$. Furthermore, $z_i^h(t) \rightarrow x_i(t)$ uniformly as $h \rightarrow 0$, and

$$|x_i(t) - z_i^h(t)| \leq \frac{\beta t^\alpha e^{\beta t} w(x, h, T)}{\Gamma(\alpha + 1)}, \quad t \in [0, T]. \quad (4.26)$$

On $[\tau, T)$ the functions x_i and z_i^h are Lipschitz functions, and hence

$$w(x, h, T) \leq ch \text{ on } [\tau, T),$$

with c a positive constant. If the initial function φ is Lipschitz on $[-\tau, 0]$, then $w(x, h, T) \leq ch$ on $[0, T)$.

Corollary 1. If the initial function φ is Lipschitz on $[-\tau, 0]$, then the estimations in (4.25) and (4.26) become

$$|x_i(t) - z_i^h(t)| \leq \frac{\beta T^\alpha e^{\beta T}}{\Gamma(\alpha + 1)} ch, \quad t \in [0, T),$$

$$|x_i(nh) - z_i^h(nh)| \leq \frac{\beta e^{[\beta + \ln(\alpha)]T}}{\Gamma(\alpha + 1)} ch, \quad n \leq \left\lceil \frac{T}{h} \right\rceil$$

and for $\alpha \rightarrow 1$ the result is also true.

The results are true for the systems purely ordinary and purely delayed.

Corollary 2. The assertion of Theorem 4.3 remains true if $f(\cdot) \equiv 0$, i.e. the x solution of the system of differential equations

$$D_t^\alpha x_i(t) = -a_i x_i(t) + \sum_{j=1}^m c_{ij} g(x_j(t - \tau)) + I_i \quad t \in [0, T)$$

$$x_i(t) = \varphi_i(t), \quad t \in [-\tau, 0], \quad T < \infty, \quad i = 1, 2, \dots, m.$$

can be approximated using constant argument and the approximate satisfies:

$$|x_i(t) - z_i^h(t)| \leq \frac{\beta_1 T^\alpha e^{\beta_1 T}}{\Gamma(\alpha + 1)} ch, \quad t \in [0, T),$$

$$|x_i(nh) - z_i^h(nh)| \leq \frac{\beta_1 e^{[\beta_1 + \ln(\alpha)]T}}{\Gamma(\alpha + 1)} ch, \quad n \leq \left\lceil \frac{T}{h} \right\rceil$$

and for $\alpha \rightarrow 1$ the result is also true, with $\beta_1 = \sum_{i=1}^m a_i + \sum_{1 \leq i, j \leq m} |c_{ij}| L_g$.

Corollary 3. The assertion of Theorem 4.3 remains true if $g(\cdot) \equiv 0$, i.e. the x solution of the system of differential equations

$$D_t^\alpha x_i(t) = -a_i x_i(t) + \sum_{j=1}^m b_{ij} f(x_j(t)) + I_i \quad t \in [0, T]$$

$$x_i(0) = x_i^0, \quad i = 1, 2, \dots, m.$$

can be approximated using constant argument and the approximate satisfies:

$$|x_i(t) - z_i^h(t)| \leq \frac{\beta_2 T^\alpha e^{\beta_2 T}}{\Gamma(\alpha + 1)} ch, \quad t \in [0, T],$$

$$|x_i(nh) - z_i^h(nh)| \leq \frac{\beta_2 e^{[\beta_2 + \ln(\alpha)]T}}{\Gamma(\alpha + 1)} ch, \quad n \leq \left\lceil \frac{T}{h} \right\rceil$$

and for $\alpha \rightarrow 1$ the result is also true, with $\beta_2 = \sum_{i=1}^m a_i + \sum_{1 \leq i, j \leq m} |b_{ij}| L_f$.

5 Numerical Result

In this section we present two numerical examples to validate the obtained theoretical results. However, before to present the details we note that the sequence \mathbf{a}_i^h in Theorem 4.1 can be rewritten. Indeed, we have that if $\mathbf{a}_i^h(n) = z_i^h(nh)$, then the sequences satisfy the following difference equations

$$z_i^h(t) = \mathbf{a}_i^h(n) + \frac{1}{\Gamma\alpha} \int_{nh}^t (t-s)^{\alpha-1} \left(-a_i \mathbf{a}_i^h(n) + \sum_{j=1}^m b_{ij} f(\mathbf{a}_j^h(n)) + \sum_{j=1}^m c_{ij} g(\mathbf{a}_j^h(n-\nu)) + I_i \right) ds. \tag{5.1}$$

The above problem (5.1) is a discretization of the original system (1.1).

Example 5.1. Consider the following fractional-order **bi**-directional neural network without delay:

$$D_t^\alpha x_1(t) = -x_1(t) + 2f(x_2(t)) + 5$$

$$D_t^\alpha x_2(t) = -x_2(t) - 1.5f(x_1(t)) + 10 \tag{5.2}$$

with $x_1(0) = 0.5, x_2(0) = 1$.

The activation function is $f(\cdot) = \tanh(\cdot)$.

In discrete form, we can write equation (5.2) as

$$\mathbf{a}_1^h(n+1) = \left(1 - \frac{h^\alpha}{\Gamma(\alpha + 1)} \right) \mathbf{a}_1^h(n) + \frac{h^\alpha}{\Gamma(\alpha + 1)} (2f(\mathbf{a}_2^h(n)) + 5)$$

$$\mathbf{a}_2^h(n+1) = \left(1 - \frac{h^\alpha}{\Gamma(\alpha + 1)} \right) \mathbf{a}_2^h(n) + \frac{h^\alpha}{\Gamma(\alpha + 1)} (-1.5f(\mathbf{a}_1^h(n)) + 10)$$

Actual versus Approximated solution				
t	$x_1(t)_{actual}$	$x_2(t)_{actual}$	$x_1(t)_{approx}$	$x_2(t)_{approx}$
0	0.5000	1.0000	0.5000	1.0000
0.75	3.1427	4.8282	3.9266	4.9781
1.50	5.0183	6.6172	4.7145	5.8738
2.25	5.9837	7.5345	5.1331	6.3525
3.00	5.9837	7.5345	5.4008	6.6591
3.75	6.4788	8.0050	5.5892	6.8753
5.25	6.8629	8.3698	5.8403	7.1638
6.25	6.9297	8.4332	5.9550	7.2957
8.50	6.9815	8.4824	6.1329	7.5004
10.25	6.9951	8.4954	6.2271	7.6088

For $\alpha = 0.6, h = 0.25$

Actual versus Approximated solution				
t	$x_1(t)_{actual}$	$x_2(t)_{actual}$	$x_1(t)_{approx}$	$x_2(t)_{approx}$
0	0.5000	1.0000	0.5000	1.0000
1.20	4.5899	6.1175	4.4832	5.6080
2.40	5.6167	7.1327	5.2037	6.4324
3.20	6.2061	7.7153	5.4630	6.7299
4.40	6.5444	8.0496	5.7175	7.0223
5.60	6.7385	8.2415	5.8869	7.2171
7.20	6.9139	8.4149	6.0427	7.3964
9.20	6.9716	8.4720	6.1754	7.5493
10.60	6.9837	8.4839	6.2441	7.6284
11.40	6.9906	8.4908	6.2772	7.6665

For $\alpha = 0.6, h = 0.20$

Actual versus Approximated solution				
t	$x_1(t)_{actual}$	$x_2(t)_{actual}$	$x_1(t)_{approx}$	$x_2(t)_{approx}$
0	0.5000	1.0000	0.5000	1.0000
1.10	3.4040	4.6975	4.4079	5.5167
1.50	3.4040	4.6975	4.7447	5.9030
2.20	4.4136	5.7665	5.1324	6.3484
3.10	5.1405	6.5349	5.4443	6.7071
3.70	5.1405	6.5349	5.5904	6.8751
5.20	6.0389	7.4844	5.8416	7.1643
6.20	6.3091	7.7699	5.9562	7.2963
8.50	6.6429	8.1226	6.1369	7.5046
10.30	6.8154	8.3050	6.2324	7.6146

For $\alpha = 0.6, h = 0.10$

TABLE 1. Actual versus Approximated solution for system (5.2) for $\alpha = 0.6$

t	$ \varepsilon_{x_1(t)} $	$ \varepsilon_{x_2(t)} $
0	0	0
0.75	0.7839	0.1499
1.50	0.3038	0.7434
2.25	0.8506	1.1820
3.00	0.5829	0.8754
3.75	0.8896	1.1297
5.25	1.0226	1.206
6.25	0.9747	1.1375
8.50	0.8486	0.9820
10.25	0.7680	0.8866

For $\alpha = 0.6, h = 0.25$


t	$ \varepsilon_{x_1(t)} $	$ \varepsilon_{x_2(t)} $
0	0	0
1.20	0.1067	0.5095
2.40	0.4130	0.7003
3.20	0.7431	0.9854
4.40	0.8269	1.0273
5.60	0.8516	1.0244
7.20	0.8712	1.0185
9.20	0.7962	0.9227
10.60	0.7396	0.8555
11.40	0.7134	0.8243

For $\alpha = 0.6, h = 0.20$


t	$ \varepsilon_{x_1(t)} $	$ \varepsilon_{x_2(t)} $
0	0	0
1.10	1.0039	0.8192
1.50	1.3407	1.2055
2.20	0.7188	0.5819
3.10	0.3038	0.1722
3.70	0.4499	0.3402
5.20	0.1973	0.3201
6.20	0.3529	0.4736
8.50	0.5060	0.6180
10.30	0.5830	0.6904

For $\alpha = 0.6, h = 0.10$


TABLE 2. Error for different h for system (5.2)



(a) For $\alpha = 0.6, h = 0.25$



(b) For $\alpha = 0.6, h = 0.20$



(c) For $\alpha = 0.6, h = 0.10$

FIGURE 1. Actual versus Approximated solution for system (5.2) for various h with $\alpha = 0.6$.

The verification for results has been performed for different values of h and $\alpha = 0.6$. Table 1 and Figure 1(a)-1(c) containing solutions at different values of h and $\alpha = 0.6$ are given below. Table 2 illustrates the difference between the actual solution and approximated solution for different values of h , where absolute error $\varepsilon_i = |exact_i - approximate_i|$, $i = 1, 2$ for different values of h .

Comparing the above tables and figures, we can say that the solution of system (5.2) can be approximated through the discretized system. Since error is decreasing with increase in the value of t , thus the obtained numerical results are in support of theoretical analysis done in sub section 4.

Now, for error estimation, we proceed as follows. In this case, $a_1 = a_2 = 1, b_{11} = 0, b_{12} = 2, b_{21} = -1.5, b_{22} = 0, L_f = 1$ and $c_{ij} = 0$, for $1 \leq i, j \leq 2$. Thus for some $c > 0$, using Corollary 3, we have $\beta_2 = \sum_{i=1}^2 a_i + \sum_{1 \leq i, j \leq 2} |b_{ij}| L_f = 5.5$. Then for $t \in [0, T)$, $T < \infty$,

$$\begin{aligned} |x_1(t) - z_1^h(t)| &\leq \frac{5.5T^\alpha e^{5.5T}}{\Gamma(1 + \alpha)} ch, \\ |x_2(t) - z_2^h(t)| &\leq \frac{5.5T^\alpha e^{5.5T}}{\Gamma(1 + \alpha)} ch, \end{aligned} \quad (5.3)$$

and for $n \leq \left\lceil \frac{T}{h} \right\rceil$,

$$\begin{aligned} |x_1(nh) - z_1^h(nh)| &\leq \frac{5.5e^{[5.5+ln(\alpha)]T}}{\Gamma(1 + \alpha)} ch, \\ |x_2(nh) - z_2^h(nh)| &\leq \frac{5.5e^{[5.5+ln(\alpha)]T}}{\Gamma(1 + \alpha)} ch. \end{aligned} \quad (5.4)$$

Varying α and h for different values, we get different approximations for $x_1(t)$ and $x_2(t)$. Moreover, it is clear from (5.3) and (5.4) that $z_i^h(t) \rightarrow x_i(t)$ uniformly as $h \rightarrow 0$ supporting the theoretical analysis done in sub-section 4.2.

Example 5.2. Consider the following fractional order delayed neural network model;

$$\begin{aligned} D_t^\alpha x_1(t) &= -x_1(t) + 0.5 \tanh(x_1(t - \tau)) + 10 \\ D_t^\alpha x_2(t) &= -x_2(t) + 1.5 \tanh(x_2(t - \tau)) + 20 \end{aligned} \quad (5.5)$$

with initial conditions (10, 5) for $t \in [-\tau, 0]$. We discuss three different cases of h for $\tau = 0.05$ in this example.

Table 3 and Figures 2(a)-2(c) depict the comparison of solutions of system (5.5) for different values of h for $\alpha = 0.8$ and fixed delay $\tau = 0.05$. Table 4 illustrates the difference between actual solution and approximated solution for different values of h , where absolute error is $\varepsilon_i = |exact_i - approximate_i|$, $i = 1, 2$ for different values of h . As t is increasing, the difference between the actual solution and the approximated solution is decreasing. In other way, we can say that the approximated solution is converging to the actual solution.

For this example, error estimation proceeds as follows. In this case, $a_1 = a_2 = 1, b_{ij} = 0$, for $1 \leq i, j \leq 2$ and $c_{11} = 0.5, c_{12} = 0, c_{21} = 0, c_{22} = 1.5, L_g = 1$. Thus for some $c > 0$, using Corollary

Actual versus Approximated solution				
t	$x_1(t)_{actual}$	$x_2(t)_{actual}$	$x_1(t)_{approx}$	$x_2(t)_{approx}$
0	10.0000	5.0000	10.0000	5.0000
1.20	10.4201	19.0764	10.0929	8.0666
2.40	10.4313	19.3555	10.1327	9.3790
3.20	10.4472	19.8532	10.1686	10.5632
4.40	10.4590	20.2169	10.2010	11.6318
5.60	10.4666	20.4524	10.2302	12.5959
7.20	10.4730	20.6538	10.2803	14.2508
9.20	10.4781	20.8141	10.3211	15.5981
10.60	10.4806	20.8926	10.3386	16.1747
11.40	10.4818	20.9293	10.3544	16.6950

For $\alpha = 0.8, h = 0.05$

Actual versus Approximated solution				
t	$x_1(t)_{actual}$	$x_2(t)_{actual}$	$x_1(t)_{approx}$	$x_2(t)_{approx}$
0	10.0000	5.0000	10.0000	5.0000
1.20	10.0851	7.8083	10.0378	7.6659
2.40	10.2143	12.0720	10.1009	9.7467
3.20	10.2629	13.6769	10.1576	11.6198
4.40	10.3033	15.0084	10.2106	13.3696
5.60	10.3368	16.1132	10.2611	15.0331
7.20	10.3876	17.7909	10.3415	17.3537
9.20	10.4226	18.9460	10.3880	18.5654
10.60	10.4358	19.3807	10.4016	18.8685
11.40	10.4467	19.7414	10.4101	19.0200

For $\alpha = 0.8, h = 0.10$

Actual versus Approximated solution				
t	$x_1(t)_{actual}$	$x_2(t)_{actual}$	$x_1(t)_{approx}$	$x_2(t)_{approx}$
0	10.0000	5.0000	10.0000	5.0000
1.20	10.5280	24.4866	10.2524	13.3293
2.40	10.5209	21.6558	10.3258	15.7503
3.20	10.3976	17.8010	10.3744	17.4539
4.40	10.4403	20.0968	10.4137	18.6528
5.60	10.4983	21.5996	10.4393	19.4964
7.20	10.4604	20.2626	10.4699	20.5078
9.20	10.4838	21.0791	10.4851	21.0087
10.60	10.4787	20.8650	10.4895	21.1543
11.40	10.4792	20.9018	10.4926	21.2567

For $\alpha = 0.8, h = 0.20$

TABLE 3. Actual versus Approximated solution for $\tau = 0.05$ for system (5.5)

t	$ \epsilon_{x_1(t)} $	$ \epsilon_{x_2(t)} $
0	0	0
1.20	0.3272	11.0098
2.40	0.2986	9.9765
3.20	0.2786	9.2900
4.40	0.2580	8.5851
5.60	0.2364	7.8565
7.20	0.1927	6.4030
9.20	0.1570	5.2160
10.60	0.1420	4.7179
11.40	0.1274	4.2343

For $\alpha = 0.8, h = 0.05, \tau = 0.05$

t	$ \epsilon_{x_1(t)} $	$ \epsilon_{x_2(t)} $
0	0	0
1.20	0.0473	0.1424
2.40	0.1134	2.3253
3.20	0.1053	2.0571
4.40	0.0927	1.6388
5.60	0.0757	1.0801
7.20	0.0461	0.4372
9.20	0.0346	0.3806
10.60	0.0342	0.5122
11.40	0.0366	0.7214

For $\alpha = 0.8, h = 0.10, \tau = 0.05$

t	$ \epsilon_{x_1(t)} $	$ \epsilon_{x_2(t)} $
0	0	0
1.20	0.2756	11.1573
2.40	0.1951	5.9055
3.20	0.0232	0.3471
4.40	0.0266	1.4440
5.60	0.0590	2.1032
7.20	0.0095	0.2452
9.20	0.0013	0.0704
10.60	0.0108	0.2893
11.40	0.0134	0.3549

For $\alpha = 0.8, h = 0.20, \tau = 0.05$

TABLE 4. Error for $\alpha = 0.8, \tau = 0.05$ for system (5.5)




FIGURE 2. Actual versus Approximated solution for system (5.5) with $\alpha = 0.8, \tau = 0.05$

2, we have $\beta_1 = \sum_{i=1}^2 a_i + \sum_{1 \leq i, j \leq 2} |c_{ij}| L_g = 4$. Then for $t \in [0, T)$, $T < \infty$, we can give error estimate as

$$\begin{aligned} |x_1(t) - z_1^h(t)| &\leq ch \frac{4}{\Gamma(1+\alpha)} T^\alpha e^{4t}, \\ |x_2(t) - z_2^h(t)| &\leq ch \frac{4}{\Gamma(1+\alpha)} T^\alpha e^{4t}. \end{aligned} \quad (5.6)$$

and for $n \leq \lceil \frac{T}{h} \rceil$,

$$\begin{aligned} |x_1(nh) - z_1^h(nh)| &\leq ch \frac{4}{\Gamma(1+\alpha)} e^{(4+ln(0.8))T}, \\ |x_2(nh) - z_2^h(nh)| &\leq ch \frac{4}{\Gamma(1+\alpha)} e^{(4+ln(0.8))T}. \end{aligned} \quad (5.7)$$

In this case also, by varying h for different values, we can obtain various estimates for $x_1(t)$ and $x_2(t)$ and clearly from (5.6) and (5.7), $z_i^h(t) \rightarrow x_i(t)$ uniformly as $h \rightarrow 0$ supporting the theoretical analysis done in sub-section 4.2.

6 Conclusion

This work is devoted to study Mittag-Leffler stability for a delayed fractional-order cellular neural network containing along with the approximation of solution of model system (1.1) using discretization scheme. Later, it is proved that the solution of the original problem can be approximated for large t in a bounded interval $[0, T)$ using discretization with piecewise constant arguments of generalized type. Furthermore, the error function is estimated which establishes the uniform convergence of the fractional polygonal lines of order α as h tends to zero. Under the obtained sufficient conditions, the discrete-time analogue preserves the unique equilibrium of the continuous-time network. Moreover, the global Mittag-Leffler stability and hence global asymptotic stability indicates an improvement over the uniform stability of the continuous-time network established in [20, 21]. Hence the proposed study can be used to describe some essential aspects of fractional calculus in delayed and non-delayed neural network model system for approximating the solutions.

Acknowledgement

We are thankful to the reviewers for their valuable comments and suggestions.

S. Abbas thanks for the support of DST-SERB, IITM-FDE/SYA/14 and M. Pinto thanks for the support of FONDECYT 1120709. D. Sepúlveda thanks for the support of research project CIR 1418 at Universidad Central de Chile.

References

- [1] Sabatier, J.; Poullain, S.; Latteux, P.; Oustaloup, A. Robust speed control of a low damped electromechanical system based on CRONE control: Application to a four mass experimental

- test bench. *Nonlinear Dynamics* 38.1-4 (2004): 383-400.
- [2] Caputo, M.; Mainardi, F. A new dissipation model based on memory mechanism. *Pure and Applied Geophysics* 91.1 (1971): 134-147.
- [3] Anastasio, T.J. The fractional-order dynamics of brainstem vestibulo-oculomotor neurons. *Biological Cybernetics* 72.1 (1994): 69-79.
- [4] Mainardi, F.; Raberto, M.; Gorenflo, R.; Scalas, E. Fractional calculus and continuous-time finance II: the waiting-time distribution. *Physica A: Statistical Mechanics and its Applications* 287.3 (2000): 468-481.
- [5] Celik, V.; Yakup, D. Chaotic fractional order delayed cellular neural network. *New Trends in Nanotechnology and Fractional Calculus Applications*. Springer Netherlands (2010): 313-320.
- [6] Chen, L.; Chai, Y.; Wu, R.; Ma, T.; Zhai, H. Dynamic analysis of a class of fractional-order neural networks with delay. *Neurocomputing* 111 (2013): 190-194.
- [7] Wang, H.; Yongguang, Y.; Guoguang, W. Stability analysis of fractional-order Hopfield neural networks with time delays. *Neural Networks* 55 (2014): 98-109.
- [8] Stamova, I. Global Mittag-Leffler stability and synchronization of impulsive fractional-order neural networks with time-varying delays. *Nonlinear Dynamics* 77.4 (2014): 1251-1260.
- [9] Deng, W.; Changpin, L.; Jinhu, L. Stability analysis of linear fractional differential system with multiple time delays. *Nonlinear Dynamics* 48.4 (2007): 409-416.
- [10] Sierociuk, D.; Grzegorz, S.; Andrzej, D. Discrete fractional order artificial neural network. *Acta Mechanica et Automatica* 5 (2011): 128-132.
- [11] Hirsch, M.W. Convergent activation dynamics in continuous time networks. *Neural Networks* 2.5 (1989): 331-349.
- [12] Delavari, H.; Dumitru, B.; Jalil, S. Stability analysis of Caputo fractional-order nonlinear systems revisited. *Nonlinear Dynamics* 67.4 (2012): 2433-2439.
- [13] Guo, S.; Lihong, H. Stability of nonlinear waves in a ring of neurons with delays. *Journal of Differential Equations* 236.2 (2007): 343-374.
- [14] Zhang, S.; Yu, Y.; Wang, H. Mittag-Leffler stability of fractional-order Hopfield neural networks. *Nonlinear Analysis: Hybrid Systems* 16 (2015): 104-121.
- [15] Kilbas, A.A.; Srivastava, H.M.; Trujillo, J.J. Theory and applications of fractional differential equations. Elsevier Science Limited 204 (2006).
- [16] Miller, K.S.; Ross, B. An introduction to the fractional calculus and fractional differential equations. New York: Wiley (1993).
- [17] Podlubny, I. Fractional differential equations: an introduction to fractional derivatives, fractional differential equations, to methods of their solution and some of their applications 198. Academic Press (1998).

- [18] Henry, D. Geometric theory of Semilinear Parabolic Partial Differential Equations. Springer-Verlag, Berlin-New York (1989).
- [19] Wu, R.C.; Hei, X.D.; Chen, L.P. Finite-time stability of fractional-order neural networks with delay. *Communications in Theoretical Physics* 60 (2013): 189-193.
- [20] Yang, X.; Song, Q.; Liu, Y.; Zhao, Z. Uniform stability analysis of fractional-order BAM neural networks with delays in the leakage terms. *Abstract and Applied Analysis* (2014). Hindawi Publishing Corporation.
- [21] Chen, L.; Chai, Y.; Wu, R.; Ma, T.; Zhai, H. Dynamic analysis of a class of fractional-order neural networks with delay. *Neurocomputing* 111 (2013): 190-194.
- [22] Ke, Y.; Miao, C. Stability analysis of fractional-order CohenGrossberg neural networks with time delay. *International Journal of Computer Mathematics* 92.6 (2015): 1102-1113.
- [23] Abbas, S. Existence and attractivity of k-pseudo almost automorphic sequence solution of a model of bidirectional neural networks. *Acta Applicandae Mathematicae* 119.1 (2012): 57-74.
- [24] Abbas, S.; Xia, Y. Existence and attractivity of k-almost automorphic sequence solution of a model of cellular neural networks with delay. *Acta Mathematica Scientia* 33.1 (2013): 290-302.
- [25] Abbas, S. Pseudo almost periodic sequence solutions of discrete time cellular neural networks. *Nonlinear Analysis, Modeling and Control* 14.3 (2009): 283-301.
- [26] Chiu K.S.; Pinto, M.; Jeng J.C. Existence and global convergence of periodic solutions in recurrent neural network models with a general piecewise alternately advanced and retarded argument. *Acta Applicandae Mathematicae* 136.1 (2015): 193-216.
- [27] Pinto, M. Asymptotic equivalence of nonlinear and quasi linear differential equations with piecewise constant arguments. *Mathematical and Computer Modelling* 49 (2009): 451-458.
- [28] Pinto, M.; Robledo G. Controllability and observability for a linear time varying system with piecewise constant delay. *Acta Applicandae Mathematicae* 136.1 (2015): 193-216.
- [29] Huang, Z.; Wang, X.; Xia, Y. Exponential attractor of -almost periodic sequence solution of discrete-time bidirectional neural networks. *Simulation Modelling Practice and Theory* 18.3 (2010): 317-337.
- [30] El-Sayed, A.M.A.; Salman, S.M. On a discretization process of fractional-order Riccati differential equation. *Journal of Fractional Calculus and Applications* 4.2 (2013): 251-259
- [31] Raheem, ZF. El.; Salman, S.M. On a discretization process of fractional-order Logistic differential equation. *Journal of the Egyptian Mathematical Society* 22.3 (2014): 407-412.
- [32] Agarwal, R.P.; El-Sayed, A.M.A.; Salman, S.M. Fractional-order Chuas system: discretization, bifurcation and chaos. *Advances in Difference Equations* (2013): 320.
- [33] Ye, H.; Gao, J.; Ding, Y. A generalized Gronwall inequality and its application to a fractional differential equation. *Journal of Mathematical Analysis and Applications* 328.2 (2007): 1075-1081.