

A nonlinear nonautonomous delay differential inequality for dissipativity of Lotka-Volterra functional differential equations

Liguang Xu^{a,*}, Danhua He^b

^a Department of Applied Mathematics, Zhejiang University of Technology, Hangzhou, 310023, PR China

^b Department of Mathematics, Zhejiang International Studies University, Hangzhou, 310012, PR China

*Corresponding Author

E-mail: xlg132@126.com

Abstract

In this paper, a Lotka-Volterra functional differential equation is considered. By establishing a nonlinear nonautonomous delay differential inequality and using a generalized Barbălat's lemma, we obtain some new sufficient conditions ensuring the dissipativity of the Lotka-Volterra functional differential equation.

2000 Mathematics Subject Classification. **34K38**. 34K25.

Keywords. Lotka-Volterra functional differential equations, Nonlinear delay differential inequality, Dissipativity, Generalized Barbălat's lemma.

1 Introduction

Let H be a real or complex Hilbert space with the inner product $\langle \cdot, \cdot \rangle$ and the corresponding norm $\|\cdot\|$, X be a dense continuously imbedded subspace of H . For any given closed interval $I \subset \mathbb{R}$, let the symbol $C_X(I)$ denote a Banach space consisting of all continuous mappings $x : I \rightarrow X$, on which the norm is defined by $\|x\|_\infty = \max_{t \in I} \|x(t)\|$. Consider the initial value problem in Lotka-Volterra functional differential equations

$$\begin{cases} y'(t) = g(t, y(t), y(\cdot)) = \eta(t, y(t))f(t, y(t), y(\cdot)), & t \geq 0, \\ y(t) = \varphi(t), & -\tau \leq t \leq 0, \end{cases} \quad (1)$$

where τ is a positive constant, $\varphi \in C_X[-\tau, 0]$ is a given initial function, $\eta : [0, +\infty) \times X \rightarrow H$ is a nonnegative continuous function, $f : [0, +\infty) \times X \times C_X[-\tau, +\infty) \rightarrow H$, and $g : [0, +\infty) \times X \times C_X[-\tau, +\infty) \rightarrow H$ is a given locally Lipschitz continuous mapping satisfying

$$\begin{aligned} 2\Re \langle u, g(t, u, \psi(\cdot)) \rangle &\leq \eta(t, u)[\gamma(t) + \alpha(t)\|u\|^2 + \beta(t) \max_{t-\mu_2(t) \leq \theta \leq t-\mu_1(t)} \|\psi(\theta)\|^2], \\ u \in X, \psi &\in C_X[-\tau, +\infty), t \in [0, +\infty), \end{aligned} \quad (2)$$

where the functions $\mu_1(t)$ and $\mu_2(t)$ are assumed to satisfy

$$0 \leq \mu_1(t) \leq \mu_2(t) \leq t + \tau, \forall t \in [0, +\infty), \quad (3)$$

$\alpha(t)$ and $\beta(t)$ are continuous functions and $\gamma(t)$ is a bounded continuous functions on the interval $[0, +\infty)$.

Recently, Wen, Yu and Wang [1] discussed the dissipativity of (1) with $\eta(t, y(t)) \equiv 1$. They established the generalized Halanay inequality and obtained the dissipativity results of (1) with $\eta(t, y(t)) \equiv 1$. In this paper, we will improve the inequality in [1] such that it is effective for (1). By establishing a nonlinear nonautonomous delay differential inequality and using a generalized Barbălat's lemma, we obtain some new sufficient conditions ensuring the dissipativity of (1).

2 Nonlinear delay differential inequality

Theorem 2.1. *If $y(t) \geq 0$ is a differentiable function defined on $(-\infty, +\infty)$, and*

$$\begin{cases} u'(t) \leq \eta(t, y(t))[\gamma(t) + \alpha(t)u(t) + \beta(t) \sup_{t-\tau(t) \leq \theta \leq t} u(\theta)], t \geq t_0, \\ y(t) = \psi(t), t \leq t_0, \end{cases} \quad (4)$$

where $u(t) = \|y(t)\|^2$, and $\psi(t)$ is bounded and continuous for $t \leq t_0$, continuous functions $\gamma(t) \geq 0$, $\beta(t) \geq 0$ and $\alpha(t) \leq 0$ for $t \in [t_0, +\infty)$, $\tau(t) \geq 0$, $\eta : [0, +\infty) \times X \rightarrow H$ is a nonnegative continuous function, and if there exists $\sigma > 0$ such that

$$\alpha(t) + \beta(t) \leq -\sigma < 0 \text{ for } t \geq t_0, \quad (5)$$

then we have

(i)

$$u(t) \leq \frac{\gamma^*}{\sigma} + G, t \geq t_0. \quad (6)$$

(ii)

$$u(t) \leq \frac{\gamma^*}{\sigma} + Ge^{-\mu^* \int_{t_0}^t \eta(s, y(s)) ds}, t \geq t_0, \quad (7)$$

where $G = \sup_{-\infty < \theta \leq t_0} \|\psi(\theta)\|^2$, $\gamma^* = \sup_{t_0 \leq t < +\infty} \gamma(t)$, and $\mu^* \geq 0$ is defined as

$$\mu^* = \inf_{t \geq t_0} \{\mu(t) : \mu(t) + \alpha(t) + \beta(t)e^{h\mu(t)\tau(t)} = 0\}, \quad (8)$$

where

$$h = \sup_{t \geq t_0} \max_{(s, u) \in [t-\tau, t] \times [0, \frac{\gamma^*}{\sigma} + G]} \eta(t, y) < \infty. \quad (9)$$

Proof. (i): We at first shall prove that for any positive constant ε ,

$$u(t) \leq \frac{\gamma^* + \varepsilon}{\sigma} + G, t \geq t_0. \quad (10)$$

If (10) does not hold, then there exists $t_1 > t_0$ such that

$$u(t_1) = \frac{\gamma^* + \varepsilon}{\sigma} + G, u'(t_1) > 0, u(t) \leq \frac{\gamma^* + \varepsilon}{\sigma} + G, t \in (-\infty, t_1]. \quad (11)$$

Using (4), (5) and (11), we obtain that

$$\begin{aligned}
 u'(t_1) &\leq \eta(t_1, y(t_1))[\gamma(t_1) + \alpha(t_1)u(t_1) + \beta(t_1) \sup_{t_1 - \tau(t_1) \leq \theta \leq t_1} u(\theta)] \\
 &\leq \eta(t_1, y(t_1))[\gamma^* + \alpha(t_1)(\frac{\gamma^* + \varepsilon}{\sigma} + G) + \beta(t_1)(\frac{\gamma^* + \varepsilon}{\sigma} + G)] \\
 &\leq \eta(t_1, y(t_1))[\frac{\gamma^* + \varepsilon}{\sigma}(\sigma + \alpha(t_1) + \beta(t_1)) + G(\alpha(t_1) + \beta(t_1))] \\
 &\leq -\eta(t_1, y(t_1))\sigma G \leq 0.
 \end{aligned} \tag{12}$$

This contradicts the inequality in (11), and so (10) holds. Since $\varepsilon > 0$ is arbitrary, we let $\varepsilon \rightarrow 0$ and obtain (6).

(ii): By (6), one can know that the definition of h for (9) is reasonable. Denote

$$H(\mu) = \mu + \alpha(t) + \beta(t)e^{h\mu\tau(t)}. \tag{13}$$

For any fixed $t \geq t_0$, we see that

$$H(0) = \alpha(t) + \beta(t) \leq -\sigma < 0, \lim_{\mu \rightarrow +\infty} H(\mu) = +\infty \tag{14}$$

and

$$H'(\mu) = 1 + \tau(t)\beta(t)he^{\mu h\tau(t)} > 0. \tag{15}$$

Therefore for any given $t \geq t_0$ there is a unique positive μ such that

$$\mu + \alpha(t) + \beta(t)e^{h\mu\tau(t)} = 0, \tag{16}$$

that means the (16) define an implicit function $\mu(t)$ for $t \geq t_0$. From that definition, one has $\mu^* \geq 0$. Next, we at first shall prove that for any positive constant ε ,

$$u(t) \leq \frac{\gamma^* + \varepsilon}{\sigma} + Ge^{-\mu^* \int_{t_0}^t \eta(s, y(s)) ds} \triangleq v(t), t \geq t_0. \tag{17}$$

If (17) is not true, then there exists a constant $\xi > t_0$ such that

$$u(\xi) = v(\xi), u'(\xi) > v'(\xi), u(t) \leq v(t), t \in [t_0, \xi]. \tag{18}$$

Let $w(t) = v(t) - u(t)$, then we have

$$\begin{aligned}
 w'(\xi) &= v'(\xi) - u'(\xi) \\
 &\geq -G\mu^* \eta(\xi, y(\xi))e^{-\mu^* \int_{t_0}^{\xi} \eta(s, y(s)) ds} - \eta(\xi, y(\xi))[\gamma(\xi) + \alpha(\xi)u(\xi) + \beta(\xi) \sup_{\xi - \tau(\xi) \leq \theta \leq \xi} u(\theta)] \\
 &> -G\mu^* \eta(\xi, y(\xi))e^{-\mu^* \int_{t_0}^{\xi} \eta(s, y(s)) ds} - \eta(\xi, y(\xi))[\gamma^* + \varepsilon + \alpha(\xi)u(\xi) + \beta(\xi) \sup_{\xi - \tau(\xi) \leq \theta \leq \xi} u(\theta)].
 \end{aligned} \tag{19}$$

If $\xi - \tau(\xi) \geq t_0$, it follows from (19) that

$$\begin{aligned}
w'(\xi) &\geq -G\mu^*\eta(\xi, y(\xi))e^{-\mu^* \int_{t_0}^{\xi} \eta(s, y(s))ds} - \eta(\xi, y(\xi))\left[\gamma^* + \varepsilon + \alpha(\xi)\left(\frac{\gamma^* + \varepsilon}{\sigma} + Ge^{-\mu^* \int_{t_0}^{\xi} \eta(s, y(s))ds}\right)\right. \\
&\quad \left. + \beta(\xi)\left(\frac{\gamma^* + \varepsilon}{\sigma} + Ge^{-\mu^* \int_{t_0}^{\xi-\tau(\xi)} \eta(s, y(s))ds}\right)\right] \\
&= \eta(\xi, y(\xi))\left[-\frac{\gamma^* + \varepsilon}{\sigma}(\sigma + \alpha(\xi) + \beta(\xi)) - Ge^{-\mu^* \int_{t_0}^{\xi} \eta(s, y(s))ds}(\mu^* + \alpha(\xi) + \beta(\xi))e^{\mu^* \int_{\xi-\tau(\xi)}^{\xi} \eta(s, y(s))ds}\right].
\end{aligned} \tag{20}$$

From the define of function $\mu(t)$, we have

$$\begin{aligned}
\mu^* + \alpha(\xi) + \beta(\xi)e^{\mu^* \int_{\xi-\tau(\xi)}^{\xi} \eta(s, y(s))ds} \\
&= \mu^* + \alpha(\xi) + \beta(\xi)e^{\mu^* \int_{\xi-\tau(\xi)}^{\xi} \eta(s, y(s))ds} - \mu(\xi) - \alpha(\xi) - \beta(\xi)e^{h\mu(\xi)\tau(\xi)} \\
&= (\mu^* - \mu(\xi)) + \beta(\xi)(e^{\mu^* \int_{\xi-\tau(\xi)}^{\xi} \eta(s, y(s))ds} - e^{h\mu(\xi)\tau(\xi)}) \leq 0.
\end{aligned} \tag{21}$$

Noting (5), therefore (20) yields

$$w'(\xi) = v'(\xi) - u'(\xi) \geq 0, \tag{22}$$

which contradicts the first inequality in (18).

If $\xi - \tau(\xi) < t_0$, it follows from (19) that

$$\begin{aligned}
w'(\xi) &\geq -G\mu^*\eta(\xi, y(\xi))e^{-\mu^* \int_{t_0}^{\xi} \eta(s, y(s))ds} - \eta(\xi, y(\xi))\left[\gamma^* + \varepsilon + \alpha(\xi)\left(\frac{\gamma^* + \varepsilon}{\sigma} + Ge^{-\mu^* \int_{t_0}^{\xi} \eta(s, y(s))ds}\right)\right. \\
&\quad \left. + \beta(\xi) \max\left\{\sup_{\theta \leq t_0} u(\theta), \sup_{t_0 \leq \theta \leq \xi} u(\theta)\right\}\right] \\
&\geq -G\mu^*\eta(\xi, y(\xi))e^{-\mu^* \int_{t_0}^{\xi} \eta(s, y(s))ds} - \eta(\xi, y(\xi))\left[\gamma^* + \varepsilon + \alpha(\xi)\left(\frac{\gamma^* + \varepsilon}{\sigma} + Ge^{-\mu^* \int_{t_0}^{\xi} \eta(s, y(s))ds}\right)\right. \\
&\quad \left. + \beta(\xi)\left(G + \frac{\gamma^* + \varepsilon}{\sigma}\right)\right] \\
&= \eta(\xi, y(\xi))\left[-\frac{\gamma^* + \varepsilon}{\sigma}(\sigma + \alpha(\xi) + \beta(\xi)) - Ge^{-\mu^* \int_{t_0}^{\xi} \eta(s, y(s))ds}(\mu^* + \alpha(\xi) + \beta(\xi))e^{\mu^* \int_{t_0}^{\xi} \eta(s, y(s))ds}\right] \\
&\geq \eta(\xi, y(\xi))\left[-\frac{\gamma^* + \varepsilon}{\sigma}(\sigma + \alpha(\xi) + \beta(\xi)) - Ge^{-\mu^* \int_{t_0}^{\xi} \eta(s, y(s))ds}(\mu^* + \alpha(\xi) + \beta(\xi))e^{\mu^* \int_{\xi-\tau(\xi)}^{\xi} \eta(s, y(s))ds}\right].
\end{aligned} \tag{23}$$

Here we also obtain that (22) holds, which contradicts the first inequality in (18). Hence the inequality (17) must hold. Since $\varepsilon > 0$ is arbitrary, we let $\varepsilon \rightarrow 0$ and obtain (7). The proof of Theorem 2.1 is completed. \square

Remark 2.2. Suppose that $\eta(t, y(t)) \equiv 1$ in Theorem 2.1, then we get Theorem 2.4 in [1].

3 Dissipativity of Lotka-Volterra functional differential equations

Definition 3.1. (See [1]) System (1) is said to be dissipative in H if there exists a bounded set $B \subset H$, such that for any given bounded set $\Phi \subset H$, there is a time $t^* = t^*(\Phi)$, such that for

any given initial function $\varphi \in C_X[-\tau, 0]$ with φ contained in Φ for all $t \in [-\tau, 0]$, the values of the corresponding solution $y(t)$ of the problem are contained in B for all $t \geq t^*$. Here B is called an absorbing set of the problem.

Lemma 3.2. (Generalized Barbălat's lemma [2]) If

- (H₁) $u : \mathbb{R}_+ \rightarrow \mathbb{R}^n$ is uniformly continuous;
- (H₂) $g : \mathbb{R}^n \rightarrow \mathbb{R}$ is continuous and $g(x) = 0$ iff $x = 0$;
- (H₃) $h : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ satisfies $\mathcal{K}(\delta) \triangleq \inf_{t \geq 0} \int_t^{t+\delta} h(s)ds > 0$, for any $\delta > 0$;
- (H₄) $\lim_{t \rightarrow \infty} \int_0^t h(s)g(u(s))ds$ exists and is finite;
then $\lim_{t \rightarrow \infty} u(t) = 0$.

Theorem 3.3. Suppose that $y(t)$ is a solution of the problem (1) satisfying the condition (2), and there exists a constant $\sigma > 0$ such that

$$\alpha(t) + \beta(t) \leq -\sigma < 0 \text{ for } t \geq 0. \tag{24}$$

Then

(i)

$$\|y(t)\|^2 \leq \frac{\bar{\gamma}^*}{\sigma} + \bar{G}, t \geq 0. \tag{25}$$

(ii)

$$\|y(t)\|^2 \leq \frac{\bar{\gamma}^*}{\sigma} + \bar{G}e^{-\mu^* \int_{t_0}^t \eta(s,y(s))ds}, t \geq 0, \tag{26}$$

where $\bar{G} = \sup_{-\infty < \theta \leq 0} \|\varphi(\theta)\|^2$, $\bar{\gamma}^* = \sup_{0 \leq t < +\infty} \gamma(t)$, and $\bar{\mu}^* \geq 0$ is defined as

$$\bar{\mu}^* = \inf_{t \geq 0} \{ \mu(t) : \mu(t) + \alpha(t) + \beta(t)e^{h\mu(t)\tau(t)} = 0 \}, \tag{27}$$

where

$$\bar{h} = \sup_{t \geq 0} \max_{(s, \|y\|^2) \in [t-\tau, t] \times [0, \frac{\bar{\gamma}^*}{\sigma} + \bar{G}]} \eta(t, y) < \infty. \tag{28}$$

Proof. To apply the result of Theorem 2.1, we have to extend the define of initial function in (1) as $y(t) = \varphi(-\tau)$ for $-\infty < t \leq -\tau$.

Let

$$u(t) = \|y(t)\|^2 = \langle y(t), y(t) \rangle. \tag{29}$$

From (2), we have

$$\begin{aligned} u'(t) &= \frac{d}{dt} \langle y(t), y(t) \rangle = 2\mathfrak{R} \langle y(t), g(t, y(t), y(\cdot)) \rangle \\ &\leq \eta(t, y(t)) [\gamma(t) + \alpha(t)u(t) + \beta(t) \max_{t-\mu_2(t) \leq \theta \leq t-\mu_1(t)} u(\theta)] \\ &\leq \eta(t, y(t)) [\gamma(t) + \alpha(t)u(t) + \beta(t) \max_{t-\mu_2(t) \leq \theta \leq t} u(\theta)]. \end{aligned} \tag{30}$$

Application of Theorem 2.1 to the above inequality yields (25) and (26). The proof is completed. \square

Corollary 3.4. *In addition to the conditions of Theorem 3.3 hold, further assume that $\eta(s, y(s)) \geq \delta > 0$. Then,*

(i) *for any given $\varepsilon > 0$, there exists a positive number $t^*(\|\varphi\|_\infty, \varepsilon)$, such that*

$$\|y(t)\|^2 \leq \frac{\bar{\gamma}^*}{\sigma} + \varepsilon, \forall t > t^*.$$

(ii) *For any given $\varepsilon > 0$, the problem (1) is dissipative with an absorbing set $B = B(0, \sqrt{\frac{\bar{\gamma}^*}{\sigma} + \varepsilon})$.*

Theorem 3.5: *In addition to the conditions of Theorem 3.3 hold, further assume that $\eta(s, y(s)) = h(s)g(y(s))$, where g and h satisfy (H_2) and (H_3) of Lemma 3.2, respectively. Then, for any given $\varepsilon > 0$, the problem (1) is dissipative with an absorbing set $B = B(0, \sqrt{\frac{\bar{\gamma}^*}{\sigma} + \varepsilon})$.*

Proof: We only need to consider the following two possible cases:

(i) If $\int_0^\infty \eta(s, y(s))ds = \infty$, then from (26) we have $\lim_{t \rightarrow \infty} \|y(t)\| \leq \sqrt{\frac{\bar{\gamma}^*}{\sigma}}$.

(ii) If $\int_0^\infty \eta(s, y(s))ds < \infty$, then $h(s)g(y(s)) \in L[0, \infty)$. From (25) and (30), we know that $\dot{y}(t)$ is bounded. So $y(t)$ is a uniformly continuous function. By Lemma 3.2, we have $\lim_{t \rightarrow \infty} y(t) = 0 \leq \sqrt{\frac{\bar{\gamma}^*}{\sigma}}$.

From above (i) and (ii), we know the problem (1) is dissipative with an absorbing set $B = B(0, \sqrt{\frac{\bar{\gamma}^*}{\sigma} + \varepsilon})$. The proof is completed. \square

Corollary 3.6. *In addition to the conditions of Theorem 3.3 hold. If $\eta(s, y(s)) = g(y(s))$, where $g(\cdot)$ is a continuous, positive definite function, then for any given $\varepsilon > 0$, system (1) is dissipative with an absorbing set $B = B(0, \sqrt{\frac{\bar{\gamma}^*}{\sigma} + \varepsilon})$.*

Remark 3.7. In the recent years, various generalized Halanay inequalities have been established and successfully applied to the problem of investigating the dissipativity of differential systems, [1,3-6]. However, the generalized Halanay inequalities in [1,3-6] are ineffective for studying the dissipativity of (1) due to the existence of the term “ $\eta(t, y(t))$ ” of (1), unless one resorts to the rather restrictive condition that $\eta(t, y(t)) \geq \delta > 0$ (δ is a constant).

Acknowledgement

The work is supported by National Natural Science Foundation of China (Grant No. 11101367) and the China Scholarship Council (Grant No. 201208330001).

References

- [1] L.P. Wen, Y.X. Yu and W.S. Wang, *Generalized Halanay inequalities for dissipativity of Volterra functional differential equations*, J. Math. Anal. Appl. 347 (2008), 169-178.

- [2] D.Y. Xu and L.G. Xu, *New results for studying a certain class of nonlinear delay differential systems*, IEEE Trans. Autom. Control 55 (2010), 1641-1645.
- [3] L.P. Wen, W.S. Wang and Y.X. Yu, *Dissipativity and asymptotic stability of nonlinear neutral delay integro-differential equations*, Nonlinear Anal. 72 (2010), 1746-1754.
- [4] H.J. Tian, *Numerical and analytic dissipativity of the h-method for delay differential equations with a bounded variable lag*, Int. J. Bifur. Chaos 14 (2004), 1839-1845.
- [5] S.Q. Gan, *Dissipativity of linear θ -methods for integro-differential equations*, Comput. Math. Appl. 52 (2006), 449-458.
- [6] S.Q. Gan, *Dissipativity of θ -methods for nonlinear Volterra delay-integro-differential equations*, J. Comput. Appl. Math. 206 (2007), 898-907.