

Categorical construction of the ring of fractions

Mitali Routaray and A. Behera

Mathematics Department, National Institute of Technology Rourkela, Rourkela - 769 008, India

E-mail: mitaray8@gmail.com, abehera@nitrkl.ac.in

Abstract

It is shown that the ring of fractions of the algebra of all bounded linear operators on a separable infinite dimensional Banach space is isomorphic to the Adams completion of the algebra with respect to a carefully chosen set of morphisms in the category of separable infinite dimensional Banach spaces and bounded linear norm preserving operators of norms at most 1.

2010 Mathematics Subject Classification. **47B07**. 55P60

Keywords. Category of fractions, Calculus of left fractions, Adams completion, Grothendieck universe, Ring of fractions.

1 Adams completion

The notion of (generalized) Adams completion arose from a general categorical completion process, suggested by Adams [1,2]. Originally, this was considered for admissible categories and generalized homology (or cohomology) theories. Subsequently, this notion has been considered in a more general framework by Deleanu, Frei and Hilton [6] where an arbitrary category and an arbitrary set of morphisms of the category are considered.

Let \mathcal{C} be a category and S be a set of morphisms of \mathcal{C} . Let $\mathcal{C}[S^{-1}]$ denote the category of fractions of \mathcal{C} with respect to S and $F : \mathcal{C} \rightarrow \mathcal{C}[S^{-1}]$, the canonical functor. Let S denote the category of sets and functions. Then for a given object Y of \mathcal{C} , $\mathcal{C}[S^{-1}](-, Y) : \mathcal{C} \rightarrow S$ defines a contravariant functor. If this functor is representable by an object Y_S of \mathcal{C} , i.e., $\mathcal{C}[S^{-1}](-, Y) \cong \mathcal{C}(-, Y_S)$, then Y_S is called the *generalized Adams completion of Y* with respect to the set of morphisms S or simply the *S -completion of Y* . We shall often refer to Y_S as the *completion of Y* [6].

The following theorem shows that under certain assumptions the Adams completion of an object of a category \mathcal{C} always exists.

1.1 Theorem. ([7], p.32, Theorem 1) *Let \mathcal{C} be a cocomplete small \mathcal{U} -category (\mathcal{U} is a fixed Grothendieck universe) and S a set of morphisms of \mathcal{C} that admits a calculus of left fractions. Suppose that the following compatibility condition with coproduct is satisfied.*

(a) *If each $s_i : X_i \rightarrow Y_i$ $i \in I$, is an element of S , where the index set I is an element of \mathcal{U} , then*

$$\bigvee_{i \in I} s_i : \bigvee_{i \in I} X_i \rightarrow \bigvee_{i \in I} Y_i$$

is an element of S .

Then every object X of \mathcal{C} has an Adams completion X_S with respect to the set of morphisms S .

The following theorem shows that under certain conditions the Adams completion of an object of a category \mathcal{C} always exists.

1.2 Theorem. ([5], p.528, Theorem 1.2) *Let S be a set of morphisms of \mathcal{C} admitting a calculus of left fractions. Then an object Y_S of \mathcal{C} is the S -completion of the object Y with respect to S if and only if there exists a morphism $e : Y \rightarrow Y_S$ in \bar{S} which is couniversal with respect to morphisms of S : given a morphism $s : Y \rightarrow Z$ in S there exists a unique morphism $t : Z \rightarrow Y_S$ in \bar{S} such that $ts = e$. In other words, the following diagram is commutative :*

$$\begin{array}{ccc} Y & \xrightarrow{e} & Y_S \\ s \downarrow & \nearrow t & \\ Z & & \end{array}$$

In order to show the morphism $e : Y \rightarrow Y_S$, as stated in Theorem 1.2 is in S , the following result is to be used.

1.3 Theorem. ([5], p.528, Theorem 1.2) *Let S be a set of morphisms in a category \mathcal{C} admitting a calculus of left fractions. Let $e : Y \rightarrow Y_S$ be the canonical morphism as defined in Theorem 1.2, where Y_S is the S -completion of Y . Furthermore, let S_1 and S_2 be sets of morphisms in the category \mathcal{C} which have the following properties:*

- (a) S_1 and S_2 are closed under composition,
- (b) $fg \in S_1$ implies that $g \in S_1$,
- (c) $fg \in S_2$ implies that $f \in S_2$,
- (d) $S = S_1 \cap S_2$.

Then $e \in S$.

2 Ring of fractions

We recall the following definitions from [10].

Let A be a ring and let S be a multiplicatively closed subset of A , i.e., $s, t \in S$ implies $st \in S$ and $1 \in S$. A *ring of fractions* (right) of A with respect to S [10] is defined as a ring $A[S^{-1}]$ together with a ring homomorphism $u : A \rightarrow A[S^{-1}]$ satisfying

- (i) $u(s)$ is invertible for every $s \in S$,
- (ii) every element in $A[S^{-1}]$ has the form $u(a)u(s)^{-1}$ with $s \in S$,
- (iii) $u(a) = 0$ if and only if $as = 0$ for some $s \in S$.

2.1 Proposition. [10] *Let S be a multiplicatively closed subset of A . Then $A[S^{-1}]$ exists, if and only if S satisfies*

- (i) if $s \in S$ and $a \in A$ then there exists $t \in S$ and $b \in A$ such that $sb = at$,
- (ii) if $sa = 0$ with $s \in S$, then $at = 0$ for some $t \in S$.

When $A[S^{-1}]$ exists, it has the form

$$A[S^{-1}] = A \times S / \sim$$

where \sim is the equivalence relation on $A \times S$ defined as $(a, s) \sim (b, t)$ if there exists $c, d \in A$ such that $ac = bd$ and $sc = td \in S$.

Addition and multiplication of $(a, s), (b, t) \in A[S^{-1}]$ are defined in the obvious way:

$$(a, s) + (b, t) = (ac + bd, u) \text{ for some } c \in A, u \text{ and } d \in S \text{ with } u = sc = td.$$

$$(a, s) \cdot (b, t) = (ac, tu) \text{ for some } c \in A \text{ and } u \in S \text{ with } sc = bu.$$

$A[S^{-1}]$ is a ring and that $a \mapsto (a, 1)$ is a ring homomorphism $u : A \rightarrow A[S^{-1}]$.

2.2 Proposition. [10] When $A[S^{-1}]$ exists, it has the following universal property: for every ring homomorphism $g : A \rightarrow B$ such that $g(s)$ is invertible in B for every $s \in S$, then there exists a unique ring homomorphism $h : A[S^{-1}] \rightarrow B$ such that $g = hu$, i.e., the following diagram is commutative:

$$\begin{array}{ccc} A & \xrightarrow{u} & A[S^{-1}] \\ g \downarrow & \swarrow h & \\ B & & \end{array}$$

2.3 Note. The objects of $A[S^{-1}]$ are of the form $u(a)u(s)^{-1}$, $a \in A$, $s \in S$. Every element of B is of the form $g(a)g(s)^{-1}$, $a \in A$, $s \in S$ and $h : A[S^{-1}] \rightarrow B$ is an isomorphism and it is unique. Thus $u : A \rightarrow A[S^{-1}]$ is surjective.

Let A be the algebra of all bounded linear operators on a separable infinite dimensional Banach space.

We prove the following.

2.4 Proposition. Let A and B be the algebras of all bounded linear operators on a separable infinite dimensional Banach space. Let $g : A \rightarrow B$ be a surjective bounded linear homomorphism such that

- (a) $g(s)$ is a unit in B ,
- (b) $g(a) = 0$ implies $as = 0$ for some $s \in S$.

Then there exists a unique ring homomorphism $\theta : B \rightarrow A[S^{-1}]$ such that $\theta g = u$, i.e., the following diagram is commutative

$$\begin{array}{ccc} A & \xrightarrow{g} & B \\ u \downarrow & \swarrow \theta & \\ A[S^{-1}] & & \end{array}$$

Proof. By Proposition 2.2 there exists a unique isomorphism $h : A[S^{-1}] \rightarrow B$ such that $g = hu$:

$$\begin{array}{ccc} A & \xrightarrow{u} & A[S^{-1}] \\ g \downarrow & \swarrow h & \\ B & & \end{array}$$

Let $\theta = h^{-1}$. for any $a \in A$, $\theta g(a) = h^{-1}g(a) = h^{-1}hu(a) = u(a)$ implying $\theta g = u$, i.e., the above diagram is commutative.

We show that θ is unique. Let there exist another $\theta' : B \rightarrow A[S^{-1}]$ such that $\theta'g = u$. For any $b \in B$, we have $\theta'(b) = \theta'(g(b')) = u(b) = \theta g(b)$, i.e., $\theta = \theta'$. This completes the proof. ■

3 The category \mathcal{B}

Let \mathcal{U} be a fixed Grothendieck universe [9]. Let \mathcal{B} denote the category of separable infinite dimensional Banach spaces and bounded linear norm preserving operators of norms at most 1. We assume that the underlying sets of the elements of \mathcal{B} are elements of \mathcal{U} . Let

$$S = \{s \text{ in } \mathcal{B} : s \text{ is a bounded linear surjective norm preserving operator of norm at most } 1\}.$$

We prove the following propositions of S .

3.1 Proposition. Let $s_i : X_i \rightarrow Y_i$ lie in S , for each $i \in I$, where the index set I is an element of \mathcal{U} . Then

$$\bigvee_{i \in I} s_i : \bigvee_{i \in I} X_i \rightarrow \bigvee_{i \in I} Y_i$$

lies in S .

Proof. Coproducts in \mathcal{B} are l_1 sums. Let $X = \bigvee_{i \in I} X_i$, where $i \in I$ and $Y = \bigvee_{i \in I} Y_i$, where $i \in I$. Define $s = \bigvee_{i \in I} s_i : X \rightarrow Y$ by the rule $s(x) = (s_i(x_i))_{i \in I}$. Clearly, s is well defined bounded linear homomorphism.

For any $(y_i)_{i \in I} \in Y$, we have $(y_i)_{i \in I} = (s_i(x_i))_{i \in I} = s(x)$ since $s_i(X_i) = Y_i$. Thus $s(X) = Y$, i.e., is surjective. That s is a bounded linear norm preserving operator of norm at most 1, can be proved easily. Hence $s = \bigvee_{i \in I} s_i$ lies in S . This completes the proof. ■

3.2 Proposition. S admits a calculus of left fractions.

Proof. Let A and B be any two objects of category \mathcal{B} . Clearly S is a closed family of morphisms of the category \mathcal{B} . We shall verify conditions (i) and (ii) of Theorem 1.3 ([6], p.67). Let $s : M \rightarrow N$ and $t : N \rightarrow P$ be two morphisms of the category \mathcal{B} . We show that if $ts \in S$ and $s \in S$, then $t \in S$. Clearly $t \in S$. Hence the condition (i) of Theorem 1.3 ([6], p.67) holds.

In order to prove condition (ii) of Theorem 1.3 ([6], p.67), consider the diagram

$$\begin{array}{ccc} A & \xrightarrow{f} & B \\ s \downarrow & & \\ C & & \end{array}$$

in \mathcal{B} with $s \in S$. We assert that the above diagram can be embedded to a weak push-out diagram

$$\begin{array}{ccc} A & \xrightarrow{f} & B \\ s \downarrow & & \downarrow t \\ C & \xrightarrow{g} & D \end{array}$$

in \mathcal{B} with $t \in S$. Let $D = (B \oplus_{l_1} C) / \bar{\Delta}$ is the quotient of the direct sum $(B \oplus_{l_1} C)$, endowed with the l_1 norm and $\bar{\Delta}$ is the closer of the subspace

$$\Delta = \{(f(a), -s(a)) : a \in A\}.$$

Define $g : C \rightarrow D$ by the rule $g(c) = (0, c) + \bar{\Delta}$ and $t : B \rightarrow D$ by the rule $t(b) = (b, 0) + \bar{\Delta}$. Clearly, the two maps are well defined and bounded linear homomorphisms. For any $a \in A, tf(a) = t(f(a)) = (f(a), 0) + \bar{\Delta} = (0, s(a)) + \bar{\Delta} = g(s(a)) = gs(a)$ implying that $tf = gs$. Hence the diagram is commutative.

In order to show that t is surjective, take an element $(b, c) + \bar{\Delta} \in D + \bar{\Delta}$. Then $(b, c) + \bar{\Delta} = (b, 0) + (0, c) + \bar{\Delta} = t(b) + g(c) = t(b) + g(s(a)) = t(b) + tf(a) = t(b + f(a))$ (since t is linear) implying t is surjective, i.e., $t \in S$.

Next let $v : B \rightarrow Z$ and $w : C \rightarrow Z$ be in category \mathcal{B} such that $vf = ws$

$$\begin{array}{ccc} A & \xrightarrow{f} & B \\ s \downarrow & & \downarrow t \\ C & \xrightarrow{g} & D \end{array} \begin{array}{c} \searrow v \\ \downarrow \\ \searrow \theta \\ \downarrow \\ \searrow w \end{array} Z$$

Define $\theta : D \rightarrow Z$, by the rule $\theta((b, c) + \bar{\Delta}) = u(b) + v(c)$. It is easy to show that θ is well defined and bounded linear homomorphism (since $\|\theta\| \leq \max \{\|u\|, \|v\|\}$). Next for any $b \in B, \theta t(b) = \theta(b, 0) = u(b)$, showing that $\theta t = v$; similarly $\theta g = w$. Thus the two triangles are commutative. This completes the proof. ■

The following result is well known.

3.3 Proposition. *The category \mathcal{B} is cocomplete.*

From Theorems 3.1, 3.2 and 3.3 we see that all the conditions of the Theorem 1.1 and Theorem 1.2 are satisfied and hence we have the following result.

3.4 Theorem. *Every object A of the category \mathcal{B} has an Adams completion A_S with respect to the set S of homomorphisms. Furthermore, there exists a homomorphism $e : A \rightarrow A_S$ in \bar{S} which is couniversal with respect to the homomorphisms in S : given a homomorphism $s : A \rightarrow B$ in S there exists a unique homomorphism $t : B \rightarrow A_S$ in \bar{S} such that $ts = e$. In other words the following diagram is commutative :*

$$\begin{array}{ccc} A & \xrightarrow{e} & A_S \\ s \downarrow & \nearrow t & \\ B & & \end{array}$$

3.5 Theorem. *The homomorphism $e : A \rightarrow A_S$ is in S .*

Proof. Let $S_1 = \{s : X \rightarrow Y \text{ in } \mathcal{B} \mid s \text{ is a surjective homomorphism}\}$ and $S_2 = \{X \rightarrow Y \text{ in } \mathcal{B} \mid s \text{ is a homomorphism}\}$. For S_1 and S_2 , it easily follows that all the conditions of Theorem 1.3 are satisfied. Therefore, $e \in S$. This completes the proof. ■

4 The result

We show that the ring of fractions $A[S^{-1}]$ of the algebra A of all bounded linear operators on a separable infinite dimensional Banach space, is precisely the Adams completion A_S of A .

4.1 Theorem. $A[S^{-1}] \cong A_S$.

Proof. Consider the diagram :


$$\begin{array}{ccc} A & \xrightarrow{u} & A[S^{-1}] \\ e \downarrow & \nearrow \varphi & \\ A_S & & \end{array}$$

By Theorem 2.4, there exists a unique homomorphism $\varphi : A_S \rightarrow A[S^{-1}]$ in S such that $\varphi e = u$. Next consider the diagram

$$\begin{array}{ccc} A & \xrightarrow{e} & A_S \\ u \downarrow & \nearrow \psi & \\ A[S^{-1}] & & \end{array}$$


By Theorem 3.4, there exists a unique homomorphism $\psi : A[S^{-1}] \rightarrow A_S$ in S such that $\psi u = e$.

In the following diagram



we have $\psi\varphi e = \psi u = e$. By the uniqueness condition of the couniversal property of e , we conclude that $\psi\varphi = 1_{A_S}$.

Also in the diagram



we have $\varphi\psi u = \varphi e = u$. By the uniqueness condition of the couniversal property of u , we conclude that $\varphi\psi = 1_{A[S^{-1}]}$. Thus $A[S^{-1}] \cong A_S$. This completes the proof. ■

It is pleasure to thank the referee for his comments which resulted in an improved presentation of the paper.

References

- [1] J. F. Adams, *Idempotent Functors in Homotopy Theory*, Manifolds, Conf. Univ. of Tokyo Press, Tokyo, (1973).
- [2] J. F. Adams, *Stable Homotopy and Generalized Homology*, The Univ. of Chicago Press, Chicago and London (1974).
- [3] J. F. Adams, *Localization and Completion*, Lecture Notes in Mathematics, Univ. of Chicago (1975).
- [4] M. F. Atiyah and I.G. MacDonalnd, *Introduction to Commutative Algebra*, Univ. of Oxford, Addison-Wesley Publ. Company (1969).
- [5] A. Behera and S. Nanda, *Mod - \mathcal{C} Postnikov approximation of a 1-connected space*, *Canad. J. Math.* 39(3) (1987), 527–543.
- [6] A. Deleanu, A. Frei and P. Hilton, *Generalized Adams Completions*, *Cahiers de Top. et Geom. Diff.* 15(1) (1974), 61–82.

- [7] A. Deleanu, *Existence of the Adams completions for objects of cocomplete categories*, J. Pure and Appl. Alg. 6 (1975), 31–39.
- [8] P. Gabriel and M. Zisman, *Calculus of fractions and homotopy theory*, Springer Verlag, New York (1967).
- [9] H. Schubert, *Categories*, Springer-Verlag, New York (1972).
- [10] Bo Stenstrom, *Rings of Quotients*, Springer-Verlag, Berlin Heidelberg New York (1975).