

MOD 2 MORAVA  $K$ -THEORY FOR FROBENIUS  
COMPLEMENTS OF EXPONENT DIVIDING  $2^n \cdot 9$

MALKHAZ BAKURADZE

(communicated by Vladimir Vershinin)

*Abstract*

We determine the cohomology rings  $K(s)^*(B\mathcal{G})$  at 2 for all finite Frobenius complements  $\mathcal{G}$  of exponent dividing  $2^n \cdot 9$ .

Let  $V$  be an abelian group, and let  $\mathcal{G}$  be a group of automorphisms of  $V$ . If  $\mathcal{G}$  has exponent  $2^n \cdot 3^k$  for  $0 \leq n$  and  $0 \leq k \leq 2$  and  $\mathcal{G}$  acts freely on  $V$ , then  $\mathcal{G}$  is finite (see [6] Theorem 1.1). Every finite group that acts freely on an abelian group is isomorphic to a Frobenius complement in some finite Frobenius group (see [6] Lemma 2.6). By the classification of finite Frobenius complements (see [7]) the quotient of  $\mathcal{G}$  by its maximal normal 3-subgroup  $\mathcal{H}$  is isomorphic to a cyclic 2-group  $\mathcal{C}$ , a generalized quaternion group  $Q$ , the binary tetrahedral group  $2\mathcal{T}$  of order 24 (or  $\mathrm{SL}(2,3)$ ), or the binary octahedral group  $2\mathcal{O}$  of order 48. Then Atiyah-Hirzebruch-Serre spectral sequence for  $\mathcal{H} \triangleleft \mathcal{G}$  implies that at 2 the ring  $K(s)^*(B\mathcal{G})$  is isomorphic to  $K(s)^*(B\mathcal{K})$ , for  $\mathcal{K} = \mathcal{G}/\mathcal{H}$  is either  $\mathcal{C}, Q, 2\mathcal{T}, 2\mathcal{O}$ . For the cyclic group  $\mathcal{C} = \mathbb{Z}/2^k$ ,  $K(s)^*(B\mathbb{Z}/2^k) = \mathbb{F}_2[v_s, v_s^{-1}][u]/(u^{2^{ks}})$ . For the generalized quaternion group  $Q_{2^{m+2}}$  we have Theorem 1.1 of [4]. We deduce Morava  $K$ -theory rings at 2 for the groups  $2\mathcal{T}$  and  $2\mathcal{O}$  as certain subgroups in  $K(s)^*(BQ_8)$  and  $K(s)^*(BQ_{16})$  respectively (Proposition 5 and Proposition 6.)

In [3] we proved the following formula for the first Chern class of the transferred line complex bundle: Let  $X \rightarrow Y$  be the regular two covering defined by free action of  $\mathbb{Z}/2$  on  $X$  and let  $\theta \rightarrow Y$  be the associated line complex bundle; Let  $\xi \rightarrow X$  be a complex line bundle and let  $\zeta \rightarrow Y$  be the plane bundle, transferred from  $\xi$  by Atiyah transfer [2]. Then for  $Tr^* : K(s)(X) \rightarrow K(s)^*(Y)$ , the transfer homomorphism [1] for our covering  $X \rightarrow Y$ , one has

$$Tr^*(c_1(\xi)) = c_1(\theta) + c_1(\zeta) + v_s \sum_{i=1}^{s-1} c_1(\theta)^{2^s - 2^i} c_2(\zeta)^{2^{i-1}}. \quad (1)$$

We show that formula 1 plays major role in the ring structure  $K(s)^*(B\mathcal{G})$  at 2 for aforementioned groups and gives another derivations for some related rank one Lie groups.

Much of our note is written in terms of Theorem 1.1 of [4]. Let

$$G = \langle a, b \mid a^{2^{m+1}} = 1, b^2 = a^e, bab^{-1} = a^r \rangle, \quad m \geq 1$$

and either  $e = 0, r = -1$  (the dihedral group  $D_{2^{m+2}}$  of order  $2^{m+2}$ ),  $e = 2^m, r = -1$  (the generalized quaternion group  $Q_{2^{m+2}}$ ) or  $m \geq 2, e = 0, r = 2^m - 1$  (the semidihedral group  $SD_{2^{m+2}}$ ).

Spectral sequence consideration (see [8]) imply that  $K(s)(BG)$  is generated by following Chern classes  $|c| = |x| = 2, |c_2| = 4$ :

$$\begin{aligned} c &= c_1(\eta_1), \quad \eta_1 : G/\langle a \rangle \cong \mathbf{Z}/2 \rightarrow \mathbb{C}^*, \quad b \mapsto -1; \\ x &= c_1(\eta_2), \quad \eta_2 : G/\langle a^2, b \rangle \cong \mathbf{Z}/2 \rightarrow \mathbb{C}^*, \quad a \mapsto -1; \end{aligned}$$

and  $c_2 = c_2(\xi_{\pi_1})$ , where  $\xi_{\pi_1} \rightarrow B\langle a, b \rangle$  is the plane bundle transferred from the canonical line bundle  $\xi \rightarrow B\langle a \rangle$ , for the double covering  $\pi_1 : B\langle a \rangle \rightarrow B\langle a, b \rangle$  corresponding to  $\eta_1$ .

The ring structure is the result of the formula for transferred first Chern class 1. See [4].

Let  $N$  be the normalizer of  $U(1)$  in  $S^3$ . The normalizes of the maximal torus in  $SO(3)$  is  $O(2) = U(1) \rtimes \mathbf{Z}/2$  and  $\mathbf{Z}/2$  acts on  $K(s)^*BU(1) = K(s)^*[u]$  by  $[-1]_F(u)$  as above.

Since  $BU(1)_p = [\text{colim}_n B\mathbf{Z}/(p^n)]_p$ , we have

$$K(s)^*(BO(2)) = K(s)^*(\text{lim}_m(BD_{2^{m+2}})) = K(s)^*(\text{lim}_m(BSD_{2^{m+2}}))$$

and

$$K(s)^*(BN) = K(s)^*(\text{lim}_m(BQ_{2^{m+2}})).$$

Thus Theorem 1.1 of [4] implies

**Corollary 1.**  $K(s)^*(BO(2)) = K(s)^*[[c, c_2]]/(c^{2^s}, v_s c \sum_{i=1}^s c^{2^s-2^i} c_2^{2^{i-1}})$ , where  $c = c_1(\text{det}\eta)$  and  $c_2 = c_2(\eta)$  are the Chern classes of the bundle  $\eta \rightarrow BO(2)$ , the complexification of canonical  $O(2)$  bundle.

**Corollary 2.**  $K(s)^*(BN) = K(s)^*[[c, c_2]]/(c^{2^s}, c^2 + v_s c \sum_{i=1}^s c^{2^s-2^i} c_2^{2^{i-1}})$ , where  $c = c_1(\nu)$  is the Chern class of  $\nu$  the pullback bundle of the canonical real line bundle by  $N \rightarrow N/U(1) = \mathbf{Z}/2$  and  $c_2 = c_2(p^*(\zeta))$  is the Euler class of the pullback bundle of the canonical quaternionic line bundle by the inclusion  $N \subset S^3$ .

Then  $RP^2 \rightarrow BO(2) \rightarrow BO(3)$  is the projective bundle of the canonical  $SO(3)$  bundle. Hence the pullback of the complexification of this canonical  $SO(3)$  bundle splits over  $BO(2)$  as  $\eta \oplus \text{det}\eta$ . Note that  $c_1(\text{det}\eta) = c_1(\eta) + v_s c_2(\eta)^{2^s-1}$  modulo transfer for the covering  $BU(1) \rightarrow BO(2)$ . Thus  $K(s)^*(BSO(3))$  is subring in  $K(s)^*(BO(2))$  generated by  $v = c^2 + v_s c c_2^{2^s-1} + c_2$  and  $w = c c_2$ . This implies

**Corollary 3.**  $K(s)^*(BSO(3)) = K(s)^*[[v, w]](f_s(v, w), g_s(v, w))$ , where  $|v| = 4, |w| = 6$ , and  $f_s = f_s(v, w), g_s = g_s(v, w)$  are determined by  $f_2 = vw, g_2 = w^2$  and for  $s > 2$

$$f_s = \begin{cases} f_{s-1}^2 & s \text{ even,} \\ \frac{f_{s-1}g_{s-1}}{v} + wv^{2^{s-1}-1} & s \text{ odd,} \end{cases}$$

$$g_s = \begin{cases} g_{s-1}^2 & s \text{ odd,} \\ \frac{f_{s-1}g_{s-1}}{v} + wv^{2^{s-1}-1} & s \text{ even.} \end{cases}$$

Our main result is the following.

Let  $\mathcal{G}$  be a group acting freely on an abelian group. Let  $\mathcal{G}$  be of exponent dividing  $2^n \cdot 9$  (hence  $\mathcal{G}$  is necessarily finite, as above) and let  $\mathcal{H} \triangleleft \mathcal{G}$  be the maximal normal 3-subgroup.

**Theorem 4.** *As a ring  $K(s)^*(B\mathcal{G})$  has one of the following forms*

(i) *If  $\mathcal{G}/\mathcal{H}=Q_8$ , then  $K(s)^*(B\mathcal{G}) = K(s)^*[c, x, c_2]/R$  and the relations  $R$  are determined by*

$$c^{2^s} = x^{2^s} = 0, v_s c c_2^{2^{s-1}} = v_s \sum_{i=1}^{s-1} c^{2^s-2^i+1} c_2^{2^{i-1}} + c^2, v_s^2 c_2^{2^s} = c^2 + cx + x^2,$$

$$v_s x c_2^{2^{s-1}} = v_s \sum_{i=1}^{s-1} x^{2^s-2^i+1} c_2^{2^{i-1}} + x^2.$$

(ii) *If  $\mathcal{G}/\mathcal{H}=Q_{2^{m+2}}$ ,  $m > 1$ , then  $K(s)^*(B\mathcal{G}) = K(s)^*[c, x, c_2]/R$ , and the relations  $R$  are determined by*

$$c^{2^s} = x^{2^s} = 0, v_s c c_2^{2^{s-1}} = v_s \sum_{i=1}^{s-1} c^{2^s-2^i+1} c_2^{2^{i-1}} + c^2, v_s^{2\kappa(m)} c_2^{2^{m s}} = cx + x^2,$$

$$v_s x c_2^{2^{s-1}} = v_s x \sum_{i=1}^{s-1} c^{2^s-2^i} c_2^{2^{i-1}} + \sum_{i=1}^{m s} v_s^{1+\kappa(m)+2^{m s}-2^i} c_2^{(2^{m s}+1)2^{s-1}-(2^s-1)2^{i-1}}$$

$$+ cx,$$

where  $\kappa(m) = \frac{2^{m s}-1}{2^s-1}$ .

(iii) *If  $\mathcal{G}/\mathcal{H}=2\mathcal{T}$ , then  $K(s)^*(B\mathcal{G}) = K(s)^*[c_2]/c_2^{(2^s+1)2^{s-1}}$ .*

(iv) *If  $\mathcal{G}/\mathcal{H}=2\mathcal{O}$ , then*

$$K(s)^*(B\mathcal{G}) = K(s)^*[c, c_2]/(c^{2^s}, c^2 + v_s c \sum_{i=1}^s c^{2^s-2^i} c_2^{2^{i-1}}, c_2^{(2^s+1)2^{s-1}}).$$

(v) *If  $\mathcal{G}/\mathcal{H}=\mathbb{Z}/2^k$ , then  $K(s)^*(B\mathcal{G}) = K(s)^*[c]/c^{2^{k s}}$ .*

Here in all cases  $|c| = |x| = 2$ ,  $|c_2| = 4$ .

The statement (v) is clear. (i) and (ii) follow from Theorem 1.1 of [4] for  $Q_8$  and  $Q_{2^{m+2}}$  respectively. What remains is to consider the cases of binary tetrahedral and binary octahedral groups.

### Binary Polyhedral groups

As it is known any finite subgroup of  $SO(3)$  is either a cyclic group, a dihedral group or one of the groups of a Platonic solid: tetrahedral group  $\mathcal{T} \cong A_4$ , cube/octahedral group  $\mathcal{O} \cong S_4$ , or icosahedral group  $\mathcal{I} \cong A_5$ . We consider the preimages of the latter groups under the covering homomorphism  $S^3 \rightarrow SO(3)$ .

**Binary tetrahedral group**

Binary tetrahedral group  $2T$  as the group of 24 units in the ring of Hurwitz integers  $2T$  is given by  $\{\pm 1, \pm i, \pm j, \pm k, \frac{1}{2}(\pm 1 \pm i \pm j \pm k)\}$ .

This group can be written as a semidirect product  $2T = Q_8 \rtimes \mathbb{Z}/3$ , where  $Q_8$  is the quaternion group consisting of the 8 Lipschitz units  $\pm 1, \pm i, \pm j, \pm k$  and  $\mathbb{Z}/3$  is the cyclic group generated by  $-\frac{1}{2}(1+i+j+k)$ . The cyclic group acts on the normal subgroup  $Q_8$  by conjugation. So that the generator of  $\mathbb{Z}/3$  cyclically rotates  $i, j, k$ .

Consider now Morava  $K$ -theory at 2. Then relations of Theorem 1.1 of [4] for  $K(s)^*(BQ_8)$  imply that its subring of invariants under  $\mathbb{Z}/3$  action is generated by  $c_2$ : the generator of  $\mathbb{Z}/3$  cyclically rotates  $c, x$  and  $c + x + v_s c^{2^{s-1}} x^{2^{s-1}}$ . If ignoring the powers of  $v_s$  then the first and second elementary symmetric functions in these three symbols are equal to  $c_2^{2^{s-1}}$  and  $c_2^{2^s}$  respectively and the third is zero. It follows that  $K(s)^*(B2T) \cong [K(s)^*(BQ_8)]^{\mathbb{Z}/3}$ .

**Proposition 5.**  $K(s)^*(B2T) \cong K(s)^*[c_2]/c_2^{(2^s+1)2^{s-1}}$ , where  $|c_2| = 4$ .

**Binary octahedral group  $2O$**

This group is given as the union of the 24 Hurwitz units  $\{\pm 1, \pm i, \pm j, \pm k, \frac{1}{2}(\pm 1 \pm i \pm j \pm k)\}$  with all 24 quaternions obtained from  $\frac{1}{\sqrt{2}}(\pm 1 \pm i + 0j + 0k)$  by permutation of coordinates.

The generalized quaternion group  $Q_{16}$  forms a subgroup of  $2O$  and its conjugacy classes has 3 members. Therefore by the transfer argument  $B2O$  is a stable wedge summand of  $BQ_{16}$  after localized at 2, meaning  $K(s)^*(B2O)$  is the subring in  $K(s)^*(BQ_{16})$  at 2. We show that this is the subring generated by two symbols  $c$  and  $c_2$  of Theorem 1.1 of [4]. Namely one has

**Proposition 6.**  $K(s)^*(B2O)$  is isomorphic to

$$K(s)^*[c, c_2]/(c^{2^s}, c^2 + v_s c \sum_{i=1}^s c^{2^s-2^i} c_2^{2^{i-1}}, c_2^{(2^s+1)2^{s-1}}),$$

where  $|c| = 2, |c_2| = 4$ .

**Binary icosahedral group**

$2I$  is given as the union of the 24 Hurwitz units  $\{\pm 1, \pm i, \pm j, \pm k, \frac{1}{2}(\pm 1 \pm i \pm j \pm k)\}$  with all 96 quaternions obtained from  $\frac{1}{2}(0 \pm 1 \pm i \pm \varphi^{-1}j \pm \varphi k)$  by even permutation of coordinates. Here  $\varphi = \frac{1}{2}(1 + \sqrt{5})$  is the golden ratio. This group is isomorphic to  $SL_2(5)$ -the group of all  $2 \times 2$  matrices over  $\mathbb{F}_5$  with unit determinant.

Among other subgroups the relevant subgroup is the binary tetrahedral group formed by Hurwitz units. Then coset  $2I/2O$  has 5 members hence by the transfer argument again  $B2I$  splits off  $B2O$  after localized at 2. Thus we obtain

$$K(s)^*B(2I) \cong K(s)^*B(2T).$$

## References

- [1] J. F. Adams, *Infinite Loop Spaces*, Annals of Mathematics Studies, Princeton University Press, Princeton (1978).
- [2] M. F. Atiyah, *Characters and cohomology of finite groups*, Publ. Math. of the I.H.E.S. **9** (1961), 23–64.
- [3] M. Bakuradze and S. Priddy, *Transferred Chern classes in Morava K-theory*, Proc. Amer. Math. Soc., **132** (2004), 1855-1860.
- [4] M. Bakuradze and V. V. Vershinin, *Morava K-theory rings for the dihedral, semidihedral and generalized quaternion groups in Chern classes*, Proc. Amer. Math. Soc., **134** (2006), 3707-3714.
- [5] J. H. Conway and D. A. Smith, *On Quaternions and Octonions*, Natick, Massachusetts: AK Peters, Ltd. ISBN 1-56881-134-9, (2003).
- [6] E. Jabara and P. Mayr, *Fröbenius complements of exponent dividing  $2^m \cdot 9$* , Forum Math., **21** (2009), 217–220.
- [7] D. Passman, *Permutation groups*, W. A. Benjamin, Inc., New York-Amsterdam, 1968.
- [8] M. Tezuka and N. Yagita, *Cohomology of finite groups and Brown-Peterson cohomology*, Springer LNM **1370** (1989), 396-408.

<http://www.emis.de/ZMATH/>  
<http://www.ams.org/mathscinet>

This article may be accessed via WWW at <http://tcms.org.ge/Journals/JHRS/>

Malkhaz Bakuradze  
[malkhaz.bakuradze@tsu.ge](mailto:malkhaz.bakuradze@tsu.ge)

Tbilisi State University, Georgia