

EQUIVARIANT DOLD-THOM TOPOLOGICAL GROUPS

MARCELO A. AGUILAR AND CARLOS PRIETO

(communicated by Paul Goerss)

Abstract

Let M be a covariant coefficient system for a finite group G . In this paper we analyze several topological abelian groups, some of them new, whose homotopy groups are isomorphic to the Bredon-Illman G -equivariant homology theory with coefficients in M . We call these groups equivariant Dold-Thom topological groups and we show that they are unique up to homotopy. We use one of the new groups to prove that the Bredon-Illman homology satisfies the infinite-wedge axiom and to make some calculations of the 0th equivariant homology.

1. Introduction

Given a finite group G , a covariant coefficient system M for G , and a pointed G -set S , we defined in [2] an abelian group $F^G(S, M)$. Using this construction, we associate to a pointed G -space X a simplicial abelian group $F^G(\mathcal{S}(X), M)$. Its geometric realization, denoted by $F^G(X, M)$, is a topological group whose homotopy groups are isomorphic to the reduced Bredon-Illman G -equivariant homology of X with coefficients in M . Given a G -equivariant ordinary covering map $p : E \rightarrow X$, we also defined a continuous transfer $t_p^G : F^G(X^+, M) \rightarrow F^G(E^+, M)$, that induces a transfer in equivariant homology, when M is a Mackey functor.

In [3] we showed that when the G -space X is a strong ρ -space (e.g. a G -simplicial complex or a finite dimensional countable locally finite G -CW-complex), there is a topology in the abelian groups $F^G(X^\delta, M)$, where X^δ stands for the underlying set of X . With this topology, $F^G(X, M)$ is a topological group, which we denote by $\mathbb{F}^G(X, M)$. This is a smaller group than the group $F^G(X, M)$ mentioned above. It has the property that its homotopy groups are isomorphic to the reduced Bredon-Illman G -equivariant homology of X with coefficients in M , provided that M is a homological Mackey functor. Furthermore, we proved in [4] that these topological groups admit a continuous transfer for G -equivariant ramified covering maps, whose total space and base space are strong ρ -spaces.

In this paper we present a different topology for the abelian group $F^G(X^\delta, M)$ and we denote the resulting topological group by $\mathcal{F}^G(X, M)$. We prove that for any

The authors were partially supported by PAPIIT grant IN101909.

Received February 07, 2011, revised April 20, 2011; published on July 17, 2011.

2000 Mathematics Subject Classification: 55N91, 55P91, 14F43.

Key words and phrases: Equivariant homology, homotopy groups, coefficient systems, Mackey functors, topological abelian groups.

© 2011, Marcelo A. Aguilar and Carlos Prieto. Permission to copy for private use granted.

pointed G -space X of the homotopy type of a G -CW-complex, and for any coefficient system, the homotopy groups of $\mathcal{F}^G(X, M)$ are isomorphic to the reduced Bredon-Illman G -equivariant homology of X with coefficients in M . The assumptions on X and M are much weaker than those needed to define $\mathbb{F}^G(X, M)$. However, in such a generality, it does not seem possible to construct a continuous transfer even for ordinary covering G -maps.

These new topological groups $\mathcal{F}^G(X, M)$ can be used to prove the infinite wedge axiom for the Bredon-Illman homology. They also allow us to make some calculations of the 0th homology groups of a G -space X .

Those topological groups whose homotopy groups are isomorphic to the reduced Bredon-Illman G -equivariant homology of X with coefficients in M will be called *Dold-Thom topological groups*. Hence $F^G(X, M)$, $\mathbb{F}^G(X, M)$, and $\mathcal{F}^G(X, M)$ are Dold-Thom topological groups. Furthermore, all these groups are algebraically subgroups of other topological groups, so they also have another natural topology, namely the subspace topology. These topological groups will be denoted by $\overline{F}^G(X, M)$, $\overline{\mathbb{F}}^G(X, M)$, and $\overline{\mathcal{F}}^G(X, M)$. The first two were studied in [3]. In this paper we prove that these groups are isomorphic to the former, if the coefficient system M takes values on k -modules, where k is a field of characteristic 0 or a prime p that does not divide the order of G . We also show that the Dold-Thom topological groups are unique up to homotopy. Other examples of Dold-Thom topological groups were constructed by Dold and Thom [7], McCord [15], Lima-Filho [12], dos Santos [10], and Nie [17].

The paper is organized as follows. In Section 2 we give the basic definitions that are needed. Then in Section 3 we define the new topological groups $\mathcal{F}^G(X, M)$ and $\overline{\mathcal{F}}^G(X, M)$, and we prove that the former is a Dold-Thom topological group. Then in Section 4 we compare the new topological groups with the previously defined ones. In Section 5 we analyze the case of coefficients in $k\text{-Mod}$, with the field k as explained above. In Section 6 we prove the wedge axiom and we compute the 0th equivariant homology in some cases. Finally in Section 7, we study the Dold-Thom topological groups and we show that two Dold-Thom topological groups, which are locally connected and have the homotopy type of a CW-complex, are homotopy equivalent.

2. Preliminaries

We shall work in the category of k -spaces, which will be denoted by $k\text{-Top}$. We understand by a k -space a topological space X with the property that a set $W \subset X$ is closed if and only if $f^{-1}W \subset Z$ is closed for any continuous map $f : Z \rightarrow X$, where Z is any compact Hausdorff space (see [14, 18]). Given any space X , one can clearly associate to it a k -space $k(X)$ using the condition above, which is weakly homotopy equivalent to X . The product of two spaces X and Y in this category is $X \times Y = k(X \times_{\text{top}} Y)$, where $X \times_{\text{top}} Y$ is the usual topological product. Two important properties of this category are that if $p : X \rightarrow X'$ is an identification and X is a k -space, then X' is a k -space, and if $p : X \rightarrow X'$ and $q : Y \rightarrow Y'$ are identifications, then $p \times q : X \times Y \rightarrow X' \times Y'$ is an identification too. If X

is a k -space, we shall say that $A \subset X$ has the *relative k -topology* (in $k\text{-Top}$) if $A = k(A_{\text{rel}})$, where A_{rel} denotes A with the (usual) relative topology in Top . This topology is characterized by the following property: *Let Y be a k -space. Then a map $f : Y \rightarrow A$ is continuous if and only if the composite $i \circ f : Y \rightarrow X$ is continuous, where $i : A \hookrightarrow X$ is the inclusion* (see [18]).

In what follows, we shall denote by $G\text{-Top}_*$ the category of topological pointed G -spaces such that G acts trivially on the base point, or correspondingly $G\text{-}k\text{-Top}_*$. Topab will denote the category of topological abelian groups in the category of k -spaces. Recall that a *covariant coefficient system* is a covariant functor $M : \mathcal{O}(G) \rightarrow R\text{-Mod}$, where $\mathcal{O}(G)$ is the category of G -orbits G/H , $H \subset G$, and G -functions $\alpha : G/H \rightarrow G/K$, R is a commutative ring, and $R\text{-Mod}$ is the category of R -modules. A particular role will be played by the G -function $R_{g^{-1}} : G/H \rightarrow G/gHg^{-1}$, given by *right translation* by $g^{-1} \in G$, namely

$$R_{g^{-1}}(aH) = aHg^{-1} = ag^{-1}(gHg^{-1}).$$

We shall often denote aH by $[a]_H$. Observe that if X is a G -set and $x \in X$, then the canonical bijection $G/G_x \rightarrow G/G_{g_x}$ is precisely $R_{g^{-1}}$. Here G_x denotes the *isotropy subgroup* of x , namely, the maximal subgroup of G that leaves x fixed.

Let S be a pointed G -set (where the base point x_0 remains fixed under the action of G) and M a covariant coefficient system. In [2] we defined an abelian group $F(S, M)$ as follows. Consider the following union

$$\widehat{M} = \bigcup_{H \subset G} M(G/H).$$

Then

$$F(S, M) = \{u : S \rightarrow \widehat{M} \mid u(x) \in M(G/G_x), u(x_0) = 0,$$

$$\text{and } u(x) = 0 \text{ for almost every } x \in X\}.$$

Indeed, this group $F(S, M)$ is an R -module, whose structure is given by

$$(r \cdot u)(x) = ru(x) \in M(G/G_x).$$

It has as canonical generators the functions

$$lx : S \rightarrow \widehat{M} \quad \text{given by} \quad lx(x') = \begin{cases} l & \text{if } x' = x, \\ 0 & \text{if } x' \neq x, \end{cases}$$

where $x \in S$, $x \neq x_0$, and $l \in M(G/G_x)$. The group $F(S, M)$ is a functor of S as follows. Let $f : S \rightarrow T$ be a pointed G -function. Then we define $f_* : F(S, M) \rightarrow F(T, M)$ on generators by

$$f_*(lx) = M_*(\widehat{f_x})(l)f(x),$$

that is, the homomorphism whose value on y is $M_*(\widehat{f_x})(l)$ if $y = f(x)$ and 0 otherwise, where $\widehat{f_x} : G/G_x \rightarrow G/G_{f(x)}$ is the canonical surjection.

There is an action of G on $F(S, M)$ given on generators by

$$g \cdot (lx) = M_*(R_{g^{-1}})(l)(gx).$$

Then one can consider the submodule $F(S, M)^G$ of fixed points under the G -action. Let $f : S \rightarrow T$ be as above. Since f_* is clearly a G -homomorphism, it restricts to a homomorphism between the submodules of fixed points, which we denote by

$$\bar{f}_*^G : \bar{F}^G(S, M) \rightarrow \bar{F}^G(T, M).$$

This makes $\bar{F}^G(-, M)$ into a functor $G\text{-Set}_* \rightarrow R\text{-Mod}$

On the other hand, there is a surjective homomorphism $\beta_S : F(S, M) \twoheadrightarrow F(S, M)^G$ given by

$$\beta_S(lx) = \gamma_x^G(l) = \sum_{[g] \in G/G_x} M_*(R_{g^{-1}})(l)(gx),$$

that is, essentially taking the sum over the orbit of x . One can now use this to give a different functorial structure on $F(S, M)^G$, defining for a G -function $f : S \rightarrow T$ and a generator $\gamma_x^G(l)$,

$$f_*^G(\gamma_x^G(l)) = \gamma_{f(x)}^G(M_*(\hat{f}_x)(l)).$$

This defines another functor $G\text{-Set}_* \rightarrow R\text{-Mod}$, which we call $F^G(-, M)$.

These three functors are related by the commutativity of the following diagram:

$$\begin{array}{ccccc} S & & \bar{F}^G(S, M) \xrightarrow{\iota_S} & F(S, M) & \xrightarrow{\beta_S} & F^G(S, M) & (2.1) \\ \downarrow f & & \bar{f}_*^G \downarrow & \downarrow f_* & & \downarrow f_*^G & \\ T & & \bar{F}^G(T, M) \xrightarrow{\iota_T} & F(S, M) & \xrightarrow{\beta_T} & F^G(S, M). \end{array}$$

We shall use these groups to define different topological groups in the next section.

3. Topological groups and coefficient systems

First recall the following construction. Let X be a topological space and L an R -module. Then we have the R -module $F(X, L) = \{u : X \rightarrow L \mid u(x_0) = 0, \text{ and } u(x) = 0 \text{ for almost every } x \in X\}$. Following [1] we have that this R -module can be topologized as follows (see also [15]). There is a surjective function

$$\mu : \prod_k (L \times X)^k \twoheadrightarrow F(X, L)$$

given by $(l_1, x_1; \dots; l_k, x_k) \mapsto l_1x_1 + \dots + l_kx_k$. Then $F(X, L)$ has the identification topology.

Given a pointed G -space X , we denote by X^H the pointed subspace of elements of X that remain fixed under the group elements of H .

Definition 3.1. Let M be a covariant coefficient system and let X be any pointed G -space. For each subgroup $H \subset G$ consider the topological group $F(X^H, M(G/H))$ as defined above. Define $p_H : F(X^H, M(G/H)) \rightarrow F(X, M)$ by $p_H(lx) = M_*(q_{H,x})(l)$, where $x \in X^H$, $l \in M(G/H)$, and $q_{H,x} : G/H \rightarrow G/G_x$, is the

canonical projection. Now take the homomorphism

$$p_X : \prod_{H \subset G} F(X^H, M(G/H)) \rightarrow F(X, M)$$

given by $p_X((l_H x_H)_{H \subset G}) = \sum_{H \subset G} p_H(l_H x_H)$, where the product has the product topology of k -spaces. Given any generator $lx \in F(X, M)$, lx can be seen also as an element in $F(X^{G_x}, M(G/G_x))$. Therefore p_X is surjective. Give the R -module $F(X, M)$ the identification topology induced by p_X . We obtain a k -space, which we denote by $\mathcal{F}(X, M)$. By giving $\overline{F}^G(X, M)$ the relative k -topology we obtain another k -space, which we denote by $\overline{\mathcal{F}}^G(X, M)$. Moreover, by taking on $F^G(X, M)$ the identification topology given by the epimorphism β_X , we obtain another k -space, which we denote by $\mathcal{F}^G(X, M)$.

Proposition 3.2. *The groups $\mathcal{F}(X, M)$, $\overline{\mathcal{F}}^G(X, M)$, and $\mathcal{F}^G(X, M)$ are topological groups in the category of k -spaces.*

Proof. Consider the following diagram

$$\begin{array}{ccc} \prod F(X^H, M(G/H)) \times \prod F(X^H, M(G/H)) & \xrightarrow{\Pi +_H} & \prod F(X^H, M(G/H)) \\ p_X \times p_X \downarrow & & \downarrow p_X \\ \mathcal{F}(X, M) \times \mathcal{F}(X, M) & \xrightarrow{+} & \mathcal{F}(X, M) \\ \beta_X \times \beta_X \downarrow & & \downarrow \beta_X \\ \mathcal{F}^G(X, M) \times \mathcal{F}^G(X, M) & \xrightarrow{+} & \mathcal{F}^G(X, M). \end{array}$$

The function on the top is given by the product of the sum on each topological group $F(X^H, M(G/H))$ and therefore it is continuous. Since p_X and β_X are homomorphisms, both squares commute. Furthermore, since p_X and β_X are identifications, so are $p_X \times p_X$ and $\beta_X \times \beta_X$. Therefore $\mathcal{F}(X, M)$ and $\mathcal{F}^G(X, M)$ are topological groups. Finally, since $\overline{\mathcal{F}}^G(X, M)$ has the relative k -topology, it is also a topological group. \square

Proposition 3.3. *Let $f : X \rightarrow Y$ be a pointed G -map. Then the homomorphism $f_* : \mathcal{F}(X, M) \rightarrow \mathcal{F}(Y, M)$ is continuous. Thus the homomorphisms $\overline{f}_*^G : \overline{\mathcal{F}}^G(X, M) \rightarrow \overline{\mathcal{F}}^G(Y, M)$ and $f_*^G : \mathcal{F}^G(X, M) \rightarrow \mathcal{F}^G(Y, M)$ are also continuous.*

Proof. The following diagram commutes:

$$\begin{array}{ccc} \prod_{H \subset G} F(X^H, M(G/H)) & \xrightarrow{\Pi f_*^H} & \prod_{H \subset G} F(Y^H, M(G/H)) \\ p_X \downarrow & & \downarrow p_Y \\ \mathcal{F}(X, M) & \xrightarrow{f_*} & \mathcal{F}(Y, M). \end{array}$$

Indeed, if $(l_H x_H)_{H \subset G} \in \prod_{H \subset G} F(X^H, M(G/H))$ is a generator, then

$$p_Y(\prod f_*^H)((l_H x_H)_{H \subset G}) = p_Y((l_H f(x_H))_{H \subset G}) = \sum_{H \subset G} M_*(q_{f(x_H)})(l_H) f(x_H),$$

and

$$f_* p_X((l_H x_H)_{H \subset G}) = f_* \left(\sum_{H \subset G} M_*(q_{x_H})(l_H) x_H \right) = \sum_{H \subset G} M_*(\widehat{f}_{x_H}) M_*(q_{x_H})(l_H) f(x_H).$$

Here we denote q_{H, x_H} simply by q_{x_H} . Both composites are equal, since $\widehat{f}_{x_H} \circ q_{x_H} = q_{f(x_H)}$.

Since p_X is an identification and $\prod f_*^H$ is continuous, f_* is also continuous. \square

The next result shows the homotopy invariance of the functors \mathcal{F} .

Proposition 3.4. *Let $f_0, f_1 : X \rightarrow Y$ be G -homotopic pointed G -maps. Then $f_{0*}, f_{1*} : \mathcal{F}(X, M) \rightarrow \mathcal{F}(Y, M)$ are G -homotopic homomorphisms. Moreover, also $\overline{f}_{0*}^G, \overline{f}_{1*}^G : \overline{\mathcal{F}}^G(X, M) \rightarrow \overline{\mathcal{F}}^G(Y, M)$ and $f_{0*}^G, f_{1*}^G : \mathcal{F}^G(X, M) \rightarrow \mathcal{F}^G(Y, M)$ are homotopic homomorphisms.*

Proof. For each $H \subset G$, take the restriction $f_\nu^H : X^H \rightarrow Y^H$, and let $\mathcal{H} : X \times I \rightarrow Y$ be a G -homotopy such that $\mathcal{H}(x, \nu) = f_\nu(x)$, $\nu = 0, 1$, and denote by $\mathcal{H}^H : X^H \times I \rightarrow Y^H$ the restriction of \mathcal{H} , which is a homotopy between f_0^H and f_1^H . By [15], there is a homotopy $\widetilde{\mathcal{H}}^H : F(X^H, M(G/H)) \times I \rightarrow F(Y^H, M(G/H))$ between $(f_0^H)_*$ and $(f_1^H)_*$, which is given by $\widetilde{\mathcal{H}}^H(u, t) = (\mathcal{H}_t^H)_*(u)$, where $\mathcal{H}_t^H(x) = \mathcal{H}^H(x, t)$, $x \in X^H$.

Define a homotopy $R : \prod F(X^H, M(G/H)) \times I \rightarrow \prod F(Y^H, M(G/H))$ by $R((u_H)_{H \subset G}, t) = (\widetilde{\mathcal{H}}^H(u_H, t))_{H \subset G}$. Define $\mathcal{K} : \mathcal{F}(X, M) \times I \rightarrow \mathcal{F}(Y, M)$ by $\mathcal{K}(u, t) = \mathcal{H}_{t*}(u)$ and consider the diagram

$$\begin{array}{ccc} \prod F(X^H, M(G/H)) \times I & \xrightarrow{R} & \prod F(Y^H, M(G/H)) \\ p_X \times \text{id} \downarrow & & \downarrow p_Y \\ \mathcal{F}(X, M) \times I & \xrightarrow{\mathcal{K}} & \mathcal{F}(Y, M). \end{array}$$

Since both R and \mathcal{K} are homomorphisms for each fixed t , we may check the commutativity on a generator $((l_H x_H)_{H \subset G}, t) \in \prod F(X^H, M(G/H)) \times I$ as follows:

$$\begin{aligned} p_Y R((l_H x_H)_{H \subset G}, t) &= p_Y((l_H \mathcal{H}^H(x_H, t))_{H \subset G}) \\ &= \sum_{H \subset G} M_*(q_{\mathcal{H}^H(x_H, t)})(l_H) \mathcal{H}^H(x_H, t), \\ \mathcal{K}(p_X \times \text{id})((l_H x_H), t) &= \mathcal{K} \left(\sum_{H \subset G} M_*(q_{x_H})(l_H) x_H \right) \\ &= \sum_{H \subset G} M_*(\widehat{\mathcal{H}}_{t x_H}) M_*(q_{x_H})(l_H) \mathcal{H}(x_H, t). \end{aligned}$$

Both expressions are equal, since \mathcal{H}^H is the restriction of \mathcal{H} , and one clearly has that $q_{\mathcal{H}^H(x_H,t)} : G/H \rightarrow G/G_{\mathcal{H}^H(x_H,t)}$ is equal to the composite $G/H \xrightarrow{q_{x_H}} G/G_{x_H} \xrightarrow{\widehat{\mathcal{H}}_{t,x_H}} G/G_{\mathcal{H}^H(x_H,t)}$. Since $p_X \times \text{id}$ is an identification, \mathcal{K} is continuous and so it is a homotopy as desired.

The homotopy \mathcal{K} is G -equivariant. Indeed, since \mathcal{H}_{t*} is G -equivariant, one has $\mathcal{K}(g \cdot u, t) = \mathcal{H}_{t*}(g \cdot u) = g\mathcal{H}_{t*}(u) = g\mathcal{K}(u, t)$. Thus we can take the restriction of \mathcal{K} , $\overline{\mathcal{K}}^G : \overline{\mathcal{F}}^G(X, M) \times I \rightarrow \overline{\mathcal{F}}^G(Y, M)$, which is a homotopy between \overline{f}_{0*}^G and \overline{f}_{1*}^G , and by the naturality of β , we also have a commutative diagram

$$\begin{array}{ccc} \mathcal{F}(X, M) \times I & \xrightarrow{\mathcal{K}} & \mathcal{F}(Y, M) \\ \beta_X \times \text{id} \downarrow & & \downarrow \beta_Y \\ \mathcal{F}^G(X, M) \times I & \xrightarrow{\overline{\mathcal{K}}^G} & \mathcal{F}^G(Y, M), \end{array}$$

where $\mathcal{K}^G(v, t) = \mathcal{H}_{t*}^G(v)$. Thus $f_{0*}^G, f_{1*}^G : \mathcal{F}^G(X, M) \rightarrow \mathcal{F}^G(Y, M)$ are also homotopic. \square

We shall need the following G -equivariant version of the Whitehead Theorem.

Proposition 3.5. *Let Y and Z be G -spaces and $\varphi : Y \rightarrow Z$ be a G -equivariant weak homotopy equivalence. Let $y_0 \in Y$ and $\varphi(y_0) \in Z$ be base points. Assume that (Y, y_0) and $(Z, \varphi(y_0))$ have the G -homotopy type of pointed G -CW-complexes. Then $\varphi : (Y, y_0) \rightarrow (Z, \varphi(y_0))$ is a pointed G -homotopy equivalence.*

Proof. Consider the square

$$\begin{array}{ccc} (Y, y_0) & \xrightarrow{\varphi} & (Z, \varphi(y_0)) \\ \simeq_G \uparrow & & \downarrow \simeq_G \\ (C, c_0) & \xrightarrow{\psi} & (D, d_0), \end{array}$$

where (C, c_0) and (D, d_0) are pointed G -CW-complexes and ψ is defined so that the square commutes. Since the vertical arrows are G -homotopy equivalences, ψ is a G -equivariant weak homotopy equivalence.

Using [6, II(2.5)], one can show that ψ is a pointed G -homotopy equivalence. Therefore φ is also a pointed G -homotopy equivalence. \square

As a consequence of the previous two results we have the following.

Corollary 3.6. *Let Y and Z be G -spaces and $\varphi : Y \rightarrow Z$ be a G -equivariant weak homotopy equivalence. Let $y_0 \in Y$ and $\varphi(y_0) \in Z$ be base points. Assume that (Y, y_0) and $(Z, \varphi(y_0))$ have the G -homotopy type of pointed G -CW-complexes. Then φ induces a homotopy equivalence of topological groups $\varphi_*^G : \mathcal{F}^G(Y, M) \rightarrow \mathcal{F}^G(Z, M)$. \square*

Lemma 3.7. *Let (X, x_0) be a pointed G -space of the G -homotopy type of a pointed G -CW-complex, and let $\mathcal{S}(X)$ be the singular simplicial G -set of X . Let*

$\rho_X : |\mathcal{S}(X)| \rightarrow X$ be given by $\rho([\sigma, s]) = \sigma(s)$, where $\sigma : \Delta^q \rightarrow X$ and $s \in \Delta^q$. Then $\rho_{X_*}^G : \mathcal{F}^G(|\mathcal{S}(X)|, M) \rightarrow \mathcal{F}^G(X, M)$ is a homotopy equivalence of topological groups.

Proof. By [5, 1.9(e)], we have $|\mathcal{S}(X)|^H = |\mathcal{S}(X)^H|$, and clearly $|\mathcal{S}(X)^H| = |\mathcal{S}(X^H)|$. Hence $\rho_X^H : |\mathcal{S}(X)|^H \rightarrow X^H$ coincides with $\rho_{X^H} : |\mathcal{S}(X^H)| \rightarrow X^H$, which by Milnor’s theorem (see [13]) is a weak homotopy equivalence. Therefore, ρ_X is a G -equivariant weak homotopy equivalence. Hence, by the previous corollary, the result follows. \square

Definition 3.8. Assume now that Q is a simplicial pointed G -set and consider the simplicial group $F(Q, M)$ such that for any n , $F(Q, M)_n = F(Q_n, M)$. Given a morphism $\mu : \mathbf{m} \rightarrow \mathbf{n}$ in Δ , we denote by $\mu^Q : Q_n \rightarrow Q_m$ the induced pointed G -function. Then we define $\mu^{F(Q, M)} = \mu_*^Q : F(Q, M)_n \rightarrow F(Q, M)_m$. Hence one has two other simplicial groups $\overline{F}^G(Q, M)$ and $F^G(Q, M)$. $\overline{F}^G(Q, M)$ is the simplicial subgroup of $F(Q, M)$ induced by the restricted functor $\overline{F}^G(-, M)$ defined above. $F^G(Q, M)$ is the simplicial quotient group of $F(Q, M)$ induced by the quotient functor $F^G(-, M)$ defined in page 186.

Theorem 3.9. *There is a natural isomorphism of topological groups*

$$\Psi_Q : |F(Q, M)| \rightarrow \mathcal{F}(|Q|, M) \quad \text{given by} \quad \Psi_Q([lx, t]) = M_*(q_{x,t})(l)[x, t],$$

where $q_{x,t} : G/G_x \rightarrow G/G_{[x,t]}$ is the canonical projection.

Proof. In [2, 2.6] we showed that Ψ_Q is a natural isomorphism of abelian groups. Thus we only need to prove that it is a homeomorphism. First recall [5, 1.9(e)] that $|Q^H| = |Q|^H$. Now we are going to see that the following diagram commutes.

$$\begin{array}{ccc} |\prod_{H \subset G} F(Q^H, M(G/H))| & \xrightarrow{\cong} & \prod_{H \subset G} |F(Q^H, M(G/H))| \\ \downarrow |p_Q| & & \downarrow \cong \prod \psi_{Q^H} \\ & & \prod_{H \subset G} F(|Q^H|, M(G/H)) \\ & & \downarrow p_{|Q|} \\ |F(Q, M)| & \xrightarrow{\Psi_Q} & \mathcal{F}(|Q|, M). \end{array}$$

Take a generator $[(l_H x_H)_{H \subset G}, \tau] \in |\prod_{H \subset G} F(Q^H, M(G/H))|$. Then we can chase it down and we get $[\sum_{H \subset G} M_*(q_{x_H})(l_H)x_H, \tau]$, and then to the right to obtain $\sum_{H \subset G} M_*(q_{x_H, \tau})M_*(\hat{p}_{x_H})(l_H)[x_H, \tau]$. If we first go right and down with $\prod \psi_{Q^H}$, we get $(l_H[x_H, \tau])$, and then down again with $p_{|Q|}$, we obtain $\sum_{H \subset G} M_*(r_{x_H})(l_H)[x_H, \tau]$, where

$$\begin{array}{ccc} G/H & \xrightarrow{r_{x_H, \tau}} & G/G_{[x_H, \tau]} \\ & \searrow p_{x_H} & \nearrow q_{x_H, \tau} \\ & G/G_{x_H} & \end{array}$$

are the canonical projections. By [13], the isomorphism on the top is a homeomorphism, and by [1], each ψ_{Q^H} is also a homeomorphism. Since p_Q is a surjective simplicial map, its realization $|p_Q|$ is an identification. And since $p_{|Q|}$ is also an identification, then Ψ_Q is a homeomorphism. \square

Corollary 3.10. *There are natural isomorphisms of topological groups*

$$\overline{\Psi}_Q^G : |\overline{F}^G(Q, M)| \longrightarrow \overline{\mathcal{F}}^G(|Q|, M) \quad \text{and} \quad \Psi_Q^G : |F^G(Q, M)| \longrightarrow \mathcal{F}^G(|Q|, M).$$

Proof. The following is clearly a commutative diagram:

$$\begin{array}{ccccc} |\overline{F}^G(Q, M)| & \xrightarrow{\hookrightarrow} & |F(Q, M)| & \xrightarrow{|\beta_Q|} & |F^G(Q, M)| \\ \overline{\Psi}_Q^G \downarrow & & \cong \downarrow \Psi_Q & & \downarrow \Psi_Q^G \\ \overline{\mathcal{F}}^G(|Q|, M) & \xrightarrow{\hookrightarrow} & \mathcal{F}(|Q|, M) & \xrightarrow{\beta_{|Q|}} & \mathcal{F}^G(|Q|, M) \end{array}$$

and the vertical arrows on the sides are isomorphisms of abelian groups (see [2, 2.6]). Since the one in the middle, by the previous theorem, is a homeomorphism, and the first horizontal arrows are embeddings, while the second ones are identifications, $\overline{\Psi}_Q^G$ and Ψ_Q^G are homeomorphisms too. \square

The following is the main result of this section.

Theorem 3.11. *Let M be a covariant coefficient system for G and X a pointed G -space of the homotopy type of a G -CW-complex. Then the homotopy groups*

$$\pi_q(\mathcal{F}^G(X, M))$$

are naturally isomorphic to the (reduced) Bredon-Illman G -equivariant homology groups $\widetilde{H}_q^G(X; M_)$ with coefficients in M .*

Proof. $F^G(\mathcal{S}(X), M)$ is a simplicial abelian group and hence, by [13], it is a chain complex with differential $\partial_q^G : F^G(\mathcal{S}_q(X), M) \longrightarrow F^G(\mathcal{S}_{q-1}(X), M)$ given by $\partial_q^G = \sum_{i=0}^q (-1)^i (d_i^{\mathcal{S}(X)})_*^G$. By [2, 4.5] this chain complex is isomorphic to Illman’s chain complex $S^G(X, *; M)$ given in [11, p. 15], whose homology is by definition the Bredon-Illman G -equivariant reduced homology of X with coefficients in M , denoted by $\widetilde{H}_q^G(X; M)$.

We shall give an isomorphism

$$H_q(F^G(\mathcal{S}_*(X), M)) \longrightarrow \pi_q(\mathcal{F}^G(X, M)).$$

To construct the isomorphism, we shall give several isomorphisms as depicted in the following diagram.

$$\begin{array}{ccccc} H_q(F^G(\mathcal{S}(X), M)) & \xleftarrow[\cong]{i_*} & \pi_q(F^G(\mathcal{S}(X), M)) & \xrightarrow[\cong]{\Psi} & \pi_q(\mathcal{S}(|F^G(\mathcal{S}(X), M)|)) \\ \downarrow \cong & & & & \cong \downarrow \Phi \\ \pi_q(\mathcal{F}^G(X, M)) & \xleftarrow[\cong]{(\rho_*^G)_*} & \pi_q(\mathcal{F}^G(|\mathcal{S}(X)|, M)) & \xleftarrow[\cong]{\Psi_{\mathcal{S}(X)}^G} & \pi_q(|F^G(\mathcal{S}(X), M)|) \end{array}$$

By [13, 22.1], i_* is an isomorphism. In particular, this shows that every cycle in $\tilde{H}^G(X; M)$ is represented by a chain u , all of whose faces are zero. We call this a *special chain*.

The homomorphism Ψ , which is given by $\Psi(u)[\tau] = [u, \tau]$, where u is a special q -chain and $\tau \in \Delta^q$, is an isomorphism, as follows from [13, 16.6].

In order to define Φ , we must express $\Psi(u)$ as a map $\gamma : (\Delta[q], \dot{\Delta}[q]) \rightarrow (\mathcal{S}|F^H(\mathcal{S}(X), M)|, *)$. By the Yoneda lemma, γ is the unique map such that $\gamma(\delta_q) = \Psi(u)$, where $\delta_q = \text{id} : \bar{q} \rightarrow \bar{q}$. The homomorphism Φ , defined by $\Phi[\gamma][f, \tau'] = \gamma(f)(\tau')$, for $f \in \Delta[q]_n$ and $\tau' \in \Delta^n$, is given by the adjunction between the realization functor and the singular complex functor (see [13, 16.1]).

By Theorem 3.9, $\Psi_{\mathcal{S}(X)*}^G$ is an isomorphism of topological groups.

Finally, by Lemma 3.7, the homomorphism ρ_*^G is a homotopy equivalence. \square

REMARK 3.12. Chasing along the diagram and using the canonical homeomorphism $|\Delta[q]| \rightarrow \Delta^q$ given by $[f, \tau'] \mapsto f_{\#}(\tau')$, one obtains that the isomorphism $H_q(F^G(\mathcal{S}(X), M)) \rightarrow \pi_q(\mathcal{F}^G(X, M))$ is given as follows. It maps a homology class $[u]$ represented by a special chain $u = \sum_{\alpha} \gamma_{\sigma_{\alpha}}^G(u(\sigma_{\alpha}))$ to the homotopy class $[\bar{u}]$ given by $\bar{u}(\tau) = \sum_{\alpha} \gamma_{\sigma_{\alpha}(\tau)}^G(M_*(p_{\alpha})(u(\sigma_{\alpha})))$, where $p_{\alpha} : G/G_{\sigma_{\alpha}} \rightarrow G/G_{\sigma_{\alpha}(\tau)}$ is the quotient function and $\mathcal{S}(X)/G = \{[\sigma_{\alpha}]\}$.

4. Other topological abelian groups

A different way of topologizing the abelian groups $F(X^{\delta}, M)$ and $\overline{F}^G(X^{\delta}, M) = F^G(X^{\delta}, M)$, where X^{δ} denotes the underlying set of X , is as follows.

Definition 4.1. Let X be a pointed G -space (not necessarily a k -space) and M be a covariant coefficient system. Denote by $\mathcal{S}(X)$ the singular simplicial pointed G -set associated to X and consider the surjective map

$$\pi_X : |F(\mathcal{S}(X), M)| \twoheadrightarrow F(X^{\delta}, M) \quad \text{given by} \quad \pi_X([l\sigma, t]) = M_*(p_{\sigma,t})(l)\sigma(t),$$

where $p_{\sigma,t} : G/G_{\sigma} \rightarrow G/G_{\sigma(t)}$ is the canonical projection.

Give $F(X^{\delta}, M)$ the identification topology defined by π_X and denote the resulting space by $\mathbb{F}(X, M)$. Moreover, denote by $\overline{\mathbb{F}}^G(X, M)$ the group $\overline{F}^G(X^{\delta}, M)$ with the relative k -topology induced by ι_X and denote by $\mathbb{F}^G(X, M)$ the group $F^G(X^{\delta}, M)$ with the quotient topology induced by β_X .

Consider the restriction of π_X

$$\widehat{\pi}_X^G : |\overline{F}^G(\mathcal{S}(X), M)| \twoheadrightarrow \overline{F}^G(X, M)$$

and denote by $\widehat{\mathbb{F}}^G(X, M)$ the resulting identification space.

We thus have a commutative diagram

$$\begin{array}{ccccc} |\overline{F}^G(\mathcal{S}(X), M)| & \xrightarrow{|\iota_{\mathcal{S}(X)}|} & |F(\mathcal{S}(X), M)| & \xrightarrow{|\beta_{\mathcal{S}(X)}|} & |F^G(\mathcal{S}(X), M)| \\ \widehat{\pi}_X^G \downarrow & & \pi_X \downarrow & & \downarrow \pi_X^G \\ \widehat{\mathbb{F}}^G(X, M) & \xrightarrow{\text{id}} & \overline{\mathbb{F}}^G(X, M) & \xrightarrow{\iota_X} & \mathbb{F}(X, M) & \xrightarrow{\beta_X} & \mathbb{F}^G(X, M), \end{array}$$

where the vertical maps are identifications. Thus $\text{id} : \widehat{\mathbb{F}}^G(X, M) \longrightarrow \overline{\mathbb{F}}^G(X, M)$ is continuous.

REMARK 4.2. The topological groups $\overline{F}^G(X, M)$ were defined in [3].

Recall [4] that a (pointed) G -space X is called a *strong ρ -space* if the map $\rho_X : |\mathcal{S}(X)| \longrightarrow X$, given by $\rho_X([\sigma, t]) = \sigma(t)$, is a G -retraction.

Proposition 4.3. *If X is a strong ρ -space, then $\text{id} : \mathbb{F}(X, M) \longrightarrow \mathcal{F}(X, M)$ is a homeomorphism.*

Proof. Consider the diagram

$$\begin{array}{ccc} |F(\mathcal{S}(X), M)| & \xrightarrow[\cong]{\Psi_{\mathcal{S}(X)}} & \mathcal{F}(|\mathcal{S}(X)|, M) \\ \pi_X \downarrow & & \downarrow \rho_{X*} \\ \mathbb{F}(X, M) & \xrightarrow[\text{id}]{\cong} & \mathcal{F}(X, M). \end{array}$$

One easily verifies that the diagram commutes. The map on the top is a homeomorphism by 3.9, the vertical map on the left is an identification by definition, and the vertical map on the right is an identification, since it is a retraction. Thus the identity on the bottom is a homeomorphism. \square

Corollary 4.4. *If X is a strong ρ -space, then the topological groups $\mathbb{F}^G(X, M)$ and $\mathcal{F}^G(X, M)$ are equal, as well as the topological groups $\overline{\mathbb{F}}^G(X, M)$ and $\overline{\mathcal{F}}^G(X, M)$. \square*

EXAMPLE 4.5. If K is a simplicial G -set, then by [4], $|K|$ is a strong ρ -space. Hence the topological groups $\mathbb{F}(|K|, M)$ and $\mathcal{F}(|K|, M)$ are equal, and thus also the topological groups $\mathbb{F}^G(|K|, M)$ and $\mathcal{F}^G(|K|, M)$ are equal, as well as the topological groups $\overline{\mathbb{F}}^G(|K|, M)$ and $\overline{\mathcal{F}}^G(|K|, M)$.

Proposition 4.6. *If X is a strong ρ -space, then $\mathbb{F}^G(X, M) = \mathcal{F}^G(X, M)$.*

Proof. Since X is a strong ρ -space, the map $\rho_X : |\mathcal{S}(X)| \longrightarrow X$ is a G -retraction. Thus the epimorphism $\rho_X^G : \mathcal{F}^G(|\mathcal{S}(X)|, M) \twoheadrightarrow \mathcal{F}^G(X, M)$ is an identification. Consider the diagram

$$\begin{array}{ccc} |F^G(\mathcal{S}(X), M)| & \xrightarrow{\psi_M^G} & \mathcal{F}^G(|\mathcal{S}(X)|, M) \\ \pi_X^G \downarrow & & \downarrow \rho_*^G \\ \mathbb{F}^G(X, M) & \xrightarrow{\text{id}} & \mathcal{F}^G(X, M). \end{array}$$

By definition of π_X^G it commutes and by 3.9, the arrow on the top is an isomorphism of topological groups. Since both vertical arrows are identifications, the result follows. \square

There are other topological groups related to the ones previously defined, that were studied in [3].

Definition 4.7. Let X be a pointed G -space (not necessarily a k -space) and M be a covariant coefficient system for G . Define the topological group (in the category of k -spaces)

$$F(X, M) = |F(\mathcal{S}(X), M)|,$$

as well as the subgroup $\overline{F}^G(X, M) = |\overline{F}^G(\mathcal{S}(X), M)|$ and the quotient group $F^G(X, M) = |F^G(\mathcal{S}(X), M)|$. One clearly has continuous homomorphisms

$$\overline{F}^G(X, M) \xrightarrow{\iota_X} F(X, M) \xrightarrow{\beta_X} F^G(X, M)$$

given by $\iota_X = |\iota_{\mathcal{S}(X)}|$ and $\beta_X = |\beta_{\mathcal{S}(X)}|$. The first is an embedding and the second a quotient map of topological groups.

REMARK 4.8. One has identifications of topological groups

$$\overline{F}^G(X, M) \rightarrow \widehat{\mathbb{F}}^G(X, M), \quad F(X, M) \rightarrow \mathbb{F}(X, M), \quad F^G(X, M) \rightarrow \mathbb{F}^G(X, M).$$

We shall see below under which conditions the topological groups $\widehat{\mathbb{F}}^G(X, M)$ and $\overline{\mathbb{F}}^G(X, M)$ are equal.

5. Coefficients in $k\text{-Mod}$

In this section we shall assume that $M : \mathcal{O}(G) \rightarrow k\text{-Mod}$, where k is a field of characteristic 0 or a prime p that does not divide the order of G ,

Recall that if S is a pointed G -set, then we have homomorphisms

$$\overline{F}^G(S, M) \xrightarrow{\iota_S} F(S, M) \xrightarrow{\beta_S} F^G(S, M).$$

We shall study the composite $\alpha_S = \beta_S \circ \iota_S : \overline{F}^G(S, M) \rightarrow F^G(S, M)$.

Proposition 5.1. *Let S be a pointed G -set. Then α_S is a natural isomorphism.*

Proof. Take a generator

$$\gamma_x^G(l) = \sum_{[g] \in G/G_x} M_*(R_{g^{-1}})(l)(gx) \in \overline{F}^G(S, M).$$

Since $\gamma_{gx}^G M_*(R_{g^{-1}}) = \gamma_x^G$, we have

$$\begin{aligned} \alpha_S(\gamma_x^G(l)) &= \beta_S \left(\sum_{[g] \in G/G_x} M_*(R_{g^{-1}})(l)(gx) \right) \\ &= \sum_{[g] \in G/G_x} \beta_S (M_*(R_{g^{-1}})(l)(gx)) \\ &= \sum_{[g] \in G/G_x} \gamma_{gx}^G M_*(R_{g^{-1}})(l) \\ &= \sum_{[g] \in G/G_x} \gamma_x^G(l) \\ &= [G : G_x] \gamma_x^G(l) \\ &= \gamma_x^G([G : G_x]l). \end{aligned}$$

Since $p \nmid |G|$, the indexes $[G : G_x]$, seen as elements in k , are invertible elements, i.e., there exist the elements $[G : G_x]^{-1} \in k$. Since for every generator $\gamma_x^G(l)$, one has

$$\alpha_S(\gamma_x^G([G : G_x]^{-1}l)) = \gamma_x^G(l),$$

α_S is surjective. To see that it is injective, assume that $u \in \overline{F}^G(S, M)$ is such that $\alpha_S(u) = 0$. Hence

$$\alpha_S(u)(x) = [G : G_x]u(x) = 0 \quad \forall x.$$

Thus $[G : G_x]^{-1}[G : G_x]u(x) = u(x) = 0$ for all x and so $u = 0$.

The naturality of α_S follows from (2.1). □

Proposition 5.2. *Let X be a pointed G -space. Then*

- (a) $\alpha_X = |\alpha_{\mathcal{S}(X)}| : \overline{F}^G(X, M) = |\overline{F}^G(\mathcal{S}(X), M)| \longrightarrow |F^G(\mathcal{S}(X), M)| = F^G(X, M)$
and
- (b) $\alpha_X : \widehat{\mathbb{F}}^G(X, M) \longrightarrow \mathbb{F}^G(X, M)$

are isomorphisms of topological groups.

Proof. (a) follows immediately from Proposition 5.1. To see (b), consider the diagram

$$\begin{array}{ccc} |\overline{F}^G(\mathcal{S}(X), M)| & \xrightarrow{|\alpha_{\mathcal{S}(X)}|} & |F^G(\mathcal{S}(X), M)| \\ \widehat{\pi}_X^G \downarrow & & \downarrow \pi_X^G \\ \widehat{\mathbb{F}}^G(X, M) & \xrightarrow{\alpha_X} & \mathbb{F}^G(X, M). \end{array}$$

To see that it is commutative, take a generator $[\gamma_\sigma^G(l), t] \in |\overline{F}^G(\mathcal{S}(X), M)|$. Then

$$\begin{aligned} \alpha_X \widehat{\pi}_X^G([\gamma_\sigma^G(l), t]) &= \alpha_X([G_{\sigma(t)} : G_\sigma] \gamma_{\sigma(t)}^G M_*(p_\sigma)(l)) \\ &= [G_{\sigma(t)} : G_\sigma] \gamma_{\sigma(t)}^G M_*(p_\sigma)([G : G_{\sigma(t)}]l) \\ &= \gamma_{\sigma(t)}^G M_*(p_\sigma)([G : G_\sigma]l) \end{aligned}$$

and

$$\begin{aligned} \pi_X^G |_{\alpha_{\mathbb{S}(X)}}([\gamma_\sigma^G(l), t]) &= \pi_X^G([\gamma_\sigma^G([G:G_\sigma]l), t]) \\ &= \gamma_{\sigma(t)}^G M_*(p_\sigma)([G:G_\sigma]l). \end{aligned}$$

Thus we have the assertion.

Since both vertical arrows are identifications, $|\alpha_{\mathbb{S}(X)}|$ on the top is a homeomorphism and α_X on the bottom is bijective, then α_X is a homeomorphism too. \square

Corollary 5.3. *The topological groups $\widehat{\mathbb{F}}^G(X, M)$ and $\mathbb{F}^G(X, M)$ are naturally isomorphic.*

Proof. The isomorphism of topological groups α_X factors as the composite

$$\alpha_X : \widehat{\mathbb{F}}^G(X, M) \xrightarrow{\iota_X} \mathbb{F}(X, M) \xrightarrow{\beta_X} \mathbb{F}^G(X, M).$$

Therefore $\alpha_X^{-1} \circ \beta_X : \mathbb{F}(X, M) \rightarrow \widehat{\mathbb{F}}^G(X, M)$ is a left inverse for ι_X , so that ι_X is an embedding. Hence $\widehat{\mathbb{F}}^G(X, M) = \widehat{\mathbb{F}}^G(X, M) \cong \mathbb{F}^G(X, M)$. \square

6. Some applications of the homotopical Bredon-Illman homology

In this section we shall use the topological groups $\mathcal{F}^G(X, M)$ defined in Section 3, together with the G -equivariant weak homotopy equivalence axiom, that we proved in [5] using the groups $F^G(X, M)$, to show that the Bredon-Illman equivariant homology satisfies the (infinite) wedge axiom and to make some calculations. We start with a general result.

Lemma 6.1. *Let X have the homotopy type of a CW-complex. Then the connected components and the path-components of X coincide.*

Proof. Let $C : \mathcal{Top} \rightarrow \mathcal{Set}$ be the functor which associates to a space X the set $C(X)$ of its connected components. This is clearly a homotopy functor.

Now let $\varphi : X \rightarrow Y$ be a homotopy equivalence, where Y is a CW-complex. Consider the commutative diagram

$$\begin{array}{ccc} \pi_0(X) & \xrightarrow[\approx]{\varphi_*} & \pi_0(Y) \\ \downarrow & & \downarrow \\ C(X) & \xrightarrow[\approx]{C(\varphi)} & C(Y). \end{array}$$

Since Y is locally path-connected, the arrow on the right is a bijection. Hence the arrow on the left is also a bijection. \square

Definition 6.2. Let Λ be a set of indexes, and let $p(\Lambda)$ be the set of finite subsets of Λ . $p(\Lambda)$ is a directed set by inclusion. Let $\{F_\alpha \mid \alpha \in \Lambda\}$ be a family of pointed k -spaces. If $A \subset B$ are finite sets of indexes, then we have an inclusion $\prod_{\alpha \in A} F_\alpha \subset \prod_{\beta \in B} F_\beta$, defined by putting the base point $*_\beta$ in each factor F_β , whenever $\beta \notin A$. Define

$$\bigoplus_{\alpha \in \Lambda} F_\alpha = \operatorname{colim}_{A \in p(\Lambda)} \prod_{\alpha \in A} F_\alpha.$$

Since the category $k\text{-Top}$ is closed under colimits, this new space lies in $k\text{-Top}$. Notice that when $\Lambda = \mathbb{N}$ is the set of natural numbers, then the space defined above coincides with the *weak product* defined in [19].

REMARK 6.3. If the spaces F_α in the previous definition are topological abelian groups, then algebraically $\bigoplus_{\alpha \in \Lambda} F_\alpha$ is the direct sum of them, and the topology defined therein will give this sum a structure of a topological group (see next proposition). Thus, as topological groups,

$$\bigoplus_{\alpha \in \Lambda} F_\alpha = \operatorname{colim}_{A \in p(\Lambda)} \bigoplus_{\alpha \in A} F_\alpha.$$

Proposition 6.4. *If for each $\alpha \in \Lambda$, F_α is a T_1 topological group, then $\bigoplus_{\alpha \in \Lambda} F_\alpha$ is a topological group.*

Proof. Since we are taking products in $k\text{-Top}$, we consider a compact Hausdorff space K and any continuous map $f : K \rightarrow \bigoplus_{\alpha \in \Lambda} F_\alpha \times \bigoplus_{\alpha \in \Lambda} F_\alpha$. We thus have to prove that the composite

$$K \xrightarrow{f} \bigoplus_{\alpha \in \Lambda} F_\alpha \times \bigoplus_{\alpha \in \Lambda} F_\alpha \xrightarrow{+} \bigoplus_{\alpha \in \Lambda} F_\alpha$$

is continuous.

To see this, notice that the family $\{\bigoplus_{\alpha \in A} F_\alpha \mid A \in p(\Lambda)\}$ has the following two properties:

- (i) For any $A, B \in p(\Lambda)$ the index $C = A \cap B \in p(\Lambda)$ satisfies

$$\bigoplus_{\alpha \in A} F_\alpha \cap \bigoplus_{\beta \in B} F_\alpha = \bigoplus_{\gamma \in C} F_\alpha.$$

- (ii) For each $A \in p(\Lambda)$, the set $\{B \in p(\Lambda) \mid \bigoplus_{\beta \in B} F_\beta \subset \bigoplus_{\alpha \in A} F_\alpha\}$ is finite.

Therefore, by [9, 15.10], there are indexes A_1, \dots, A_m and B_1, \dots, B_n , such that $p_1 f(K) \subset \bigoplus_{\alpha \in A_1} F_\alpha \cup \dots \cup \bigoplus_{\alpha \in A_m} F_\alpha$ and $p_2 f(K) \subset \bigoplus_{\beta \in B_1} F_\beta \cup \dots \cup \bigoplus_{\beta \in B_n} F_\beta$, where p_1 and p_2 are the projections. Let $C = A_1 \cup \dots \cup A_m \cup B_1 \cup \dots \cup B_n$. Hence we have the following commutative diagram:

$$\begin{array}{ccc} & \bigoplus_{\gamma \in C} F_\gamma \times \bigoplus_{\gamma \in C} F_\gamma & \xrightarrow{+} \bigoplus_{\gamma \in C} F_\gamma \\ & \nearrow & \downarrow \\ K & \xrightarrow{f} \bigoplus_{\alpha \in \Lambda} F_\alpha \times \bigoplus_{\alpha \in \Lambda} F_\alpha & \xrightarrow{+} \bigoplus_{\alpha \in \Lambda} F_\alpha \end{array}$$

where the sum on the top is continuous because a finite product of topological groups is a topological group, and the vertical inclusion on the right is continuous. Hence the composite on the bottom is continuous. \square

Proposition 6.5. *Let X and Y be pointed spaces and L be an abelian group. Then there is an isomorphism of topological groups*

$$F(X \vee Y, L) \cong F(X, L) \times F(Y, L).$$

Proof. Consider the sequences of spaces

$$X \xrightarrow{i} X \vee Y \xrightarrow{q} Y \quad Y \xrightarrow{j} X \vee Y \xrightarrow{p} X$$

Then it follows that

$$F(X, L) \xrightarrow{i_*} F(X \vee Y, L) \xrightarrow{q_*} F(Y, L)$$

is a short exact sequence that splits. Namely, by functoriality one has that i_* is a split monomorphism, q_* is a split epimorphism and $q_* \circ i_* = 0$. Now, if $v \in F(X \vee Y, L)$ is such that $q_*(v) = 0$, then $q_*(v)(y) = v(x_0, y) = 0$. Thus $v = i_*(u)$, where $u \in F(X, L)$ is given by $u(x) = v(x, y_0)$. \square

Coming back to Definition 6.2, we have the following generalization to the infinite case of the previous proposition.

Proposition 6.6. *Let X_α , $\alpha \in \Lambda$, be a family of pointed spaces. Then there is an isomorphism of topological groups $\bigoplus_{\alpha \in \Lambda} F(X_\alpha, L) \cong F(\bigvee_{\alpha \in \Lambda} X_\alpha, L)$.*

Proof. For each $A \in p(\Lambda)$, call $\psi_A : \bigoplus_{\alpha \in A} F(X_\alpha, L) \cong F(\bigvee_{\alpha \in A} X_\alpha, L) \rightarrow F(\bigvee_{\alpha \in \Lambda} X_\alpha, L)$, where the isomorphism comes from Proposition 6.5 and the second map is induced by the canonical inclusion. If $A \subset B$, one easily verifies that $\psi_B = \psi_A \circ \psi_{A,B}$, where $\psi_{A,B} : \bigoplus_{\alpha \in A} F(X_\alpha, L) \rightarrow \bigoplus_{\alpha \in B} F(X_\alpha, L)$ is induced by the inclusion $\bigvee_{\alpha \in A} X_\alpha \subset \bigvee_{\alpha \in B} X_\alpha$ modulo the isomorphism of 6.5.

By the universal property of the colimit, the maps (continuous homomorphisms) ψ_A induce a continuous homomorphism

$$\psi : \bigoplus_{\alpha \in \Lambda} F(X_\alpha, L) \longrightarrow F(\bigvee_{\alpha \in \Lambda} X_\alpha, L).$$

In order to see that ψ is an isomorphism of topological groups, we now construct an inverse

$$\xi : F(\bigvee_{\alpha \in \Lambda} X_\alpha, L) \longrightarrow \bigoplus_{\alpha \in \Lambda} F(X_\alpha, L)$$

as follows. Let $u : \bigvee_{\alpha \in \Lambda} X_\alpha \rightarrow L$ be an element in $F(\bigvee_{\alpha \in \Lambda} X_\alpha, L)$ and let $u_\alpha : X_\alpha \rightarrow L$ be the restriction of u ; that is, $u_\alpha = p_{\alpha*}(u)$, where $p_\alpha : \bigvee_{\alpha \in \Lambda} X_\alpha \rightarrow X_\alpha$ is the canonical projection. Then $u_\alpha \neq 0$ only for finitely many values of α , i.e., only for $\alpha \in A$, and some $A \in p(\Lambda)$. Thus $(u_\alpha) \in \bigoplus_{\alpha \in \Lambda} F(X_\alpha, L)$ and one can define $\xi(u) = (u_\alpha)$. The homomorphism ξ is clearly an (algebraic) inverse of ψ . To see that ξ is continuous, define the function $\chi : L \times \bigvee_{\alpha \in \Lambda} X_\alpha \rightarrow \bigoplus_{\alpha \in \Lambda} F(X_\alpha, L)$ by

$\chi(l, x_\alpha) = lx_\alpha$. Composing χ with the identification $\prod_{\alpha \in \Lambda} L \times X_\alpha \rightarrow L \times \bigvee_{\alpha \in \Lambda} X_\alpha$ and then restricting to each $L \times X_\alpha$, we obtain the composite

$$L \times X_\alpha \xrightarrow{\mu_1} F(X_\alpha, L) \hookrightarrow \bigoplus_{\alpha \in \Lambda} F(X_\alpha, L)$$

which is continuous. Therefore χ is continuous. Now consider the commutative diagram

$$\begin{CD} (L \times \bigvee_{\alpha \in \Lambda} X_\alpha)^k @>\chi^k>> (\bigoplus_{\alpha \in \Lambda} F(X_\alpha, L))^k \\ @VVV @VV+V \\ F(\bigvee_{\alpha \in \Lambda} X_\alpha, L) @>\xi>> \bigoplus_{\alpha \in \Lambda} F(X_\alpha, L). \end{CD}$$

Since the vertical map on the left is an identification and the maps on the top and on the right are continuous (+ by 6.4), ξ is continuous. \square

Now we can use the groups $\mathcal{F}^G(X, M)$ to prove that the Bredon-Illman G -equivariant homology theory $\tilde{H}_*^G(-; M)$ satisfies the wedge axiom. We need a lemma whose proof is not difficult.

Lemma 6.7. *Let $\{Y_\alpha; i_{\beta, \alpha} : Y_\alpha \hookrightarrow Y_\beta\}$ and $\{\bar{Y}_\alpha; \bar{i}_{\beta, \alpha} : \bar{Y}_\alpha \hookrightarrow \bar{Y}_\beta\}$ be diagrams in $k\text{-Top}$ such that each $i_{\beta, \alpha}$ and $\bar{i}_{\beta, \alpha}$ are closed embeddings, and let $\{q_\alpha : Y_\alpha \rightarrow \bar{Y}_\alpha\}$ be a family of identifications such that $q_\beta \circ i_{\beta, \alpha} = \bar{i}_{\beta, \alpha} \circ q_\alpha$. Then the map $q : \text{colim } Y_\alpha \rightarrow \text{colim } \bar{Y}_\alpha$ induced in the colimits in $k\text{-Top}$ is an identification. \square*

The next lemma is a consequence of the previous one.

Lemma 6.8. *Let X_α be a pointed G -space for each $\alpha \in \Lambda$. Then the map*

$$\bigoplus_{\alpha \in \Lambda} \prod_{H \subset G} F(X_\alpha^H; M(G/H)) \xrightarrow{\oplus p_{X_\alpha}} \bigoplus_{\alpha \in \Lambda} \mathcal{F}^G(X_\alpha, M)$$

is an identification. \square

Now, as an application of our groups $\mathcal{F}^G(X, M)$, we have the next results. First we have that the Bredon-Illman homology satisfies the wedge-axiom.

Proposition 6.9. *Let $X_\alpha, \alpha \in \Lambda$, be an arbitrary family of pointed G -spaces. Then there is an isomorphism*

$$\mathcal{F}^G\left(\bigvee_{\alpha \in \Lambda} X_\alpha, M\right) \cong \bigoplus_{\alpha \in \Lambda} \mathcal{F}^G(X_\alpha, M).$$

Proof. By Proposition 6.6, the inclusions $i_\alpha : X_\alpha \hookrightarrow \bigvee_{\alpha} X_\alpha$ induce an isomorphism of topological groups which defines another isomorphism of topological groups φ such that following diagram commutes

$$\begin{CD} \bigoplus_{\alpha \in \Lambda} \prod_{H \subset G} F(X_\alpha^H; M(G/H)) @>\varphi>> \prod_{H \subset G} F(\bigvee_{\alpha \in \Lambda} X_\alpha^H, M(G/H)) \\ @V \cong VV @V \cong VV \\ \prod_{H \subset G} \bigoplus_{\alpha \in \Lambda} F(X_\alpha^H, M(G/H)) @>\psi_H>> \prod_{H \subset G} \bigoplus_{\alpha \in \Lambda} F(X_\alpha^H, M(G/H)) \end{CD}$$

Similarly, we can define an isomorphism of abelian groups

$$\psi^G : \bigoplus_{\alpha \in \Lambda} \mathcal{F}^G(X_\alpha; M) \longrightarrow \mathcal{F}^G(\bigvee_{\alpha \in \Lambda} X_\alpha, M).$$

They fit into a commutative diagram

$$\begin{array}{ccc} \bigoplus_{\alpha} \prod_{H \subset G} F(X_\alpha^H; M(G/H)) & \xrightarrow{\varphi} & \prod_{H \subset G} F(\bigvee_{\alpha} X_\alpha^H, M(G/H)) \\ \oplus p_{X_\alpha} \downarrow & & \downarrow p_{\vee X_\alpha} \\ \bigoplus_{\alpha} \mathcal{F}^G(X_\alpha, M) & \xrightarrow{\psi^G} & \mathcal{F}^G(\bigvee_{\alpha} X_\alpha, M), \end{array}$$

where the vertical arrows are identifications (the left one by the previous lemma). Since the top arrow is a homeomorphism, then the bottom arrow is a homeomorphism too. \square

Proposition 6.10. *Let $X_\alpha, \alpha \in \Lambda$, be an arbitrary family of pointed G -spaces having the homotopy type of G -CW-complexes. Then there is an isomorphism*

$$\tilde{H}_*^G(\bigvee_{\alpha} X_\alpha; M) \cong \bigoplus_{\alpha} \tilde{H}_*^G(X_\alpha; M).$$

Proof. Under conditions (i) and (ii) given in the proof of 6.4, that are satisfied by the family $\{\bigoplus_{\alpha \in A} \mathcal{F}^G(\bigvee_{\alpha} X_\alpha, M) \mid A \in p(\Lambda)\}$, it is proved in [9] that the homotopy groups commute with the colimit. Hence

$$\begin{aligned} \tilde{H}_*^G(\bigvee_{\alpha \in \Lambda} X_\alpha; M) &\cong \pi_q(\mathcal{F}^G(\bigvee_{\alpha \in \Lambda} X_\alpha; M)) && \text{(by 3.11)} \\ &\cong \pi_q(\bigoplus_{\alpha \in \Lambda} \mathcal{F}^G(X_\alpha, M)) && \text{(by 6.9)} \\ &\cong \pi_q(\text{colim}_A(\bigoplus_{\alpha \in A} \mathcal{F}^G(X_\alpha, M))) && \text{(by 6.3)} \\ &\cong \text{colim}_A(\pi_q(\bigoplus_{\alpha \in A} \mathcal{F}^G(X_\alpha, M))) && \text{(by [9, 15.9])} \\ &\cong \text{colim}_A(\bigoplus_{\alpha \in A} \pi_q(\mathcal{F}^G(X_\alpha, M))) && \text{(since } A \text{ is finite)} \\ &\cong \bigoplus_{\alpha \in \Lambda} \pi_q(\mathcal{F}^G(X_\alpha, M)) && \text{(by 6.3)} \\ &\cong \bigoplus_{\alpha \in \Lambda} \tilde{H}_*^G(X_\alpha; M) && \text{(by 3.11).} \end{aligned}$$

\square

In order to prove the wedge axiom in full generality, we need the following lemmas.

Lemma 6.11. *Let X and Y , and X' and Y' be well-pointed spaces and let $\varphi : X' \rightarrow X$ and $\psi : Y' \rightarrow Y$ be weak homotopy equivalences (mapping base points to base points). Then $\varphi \vee \psi : X' \vee Y' \rightarrow X \vee Y$ is a weak homotopy equivalence.*

Proof. Consider the double attaching spaces $X \cup I \cup Y$ and $X' \cup I \cup Y'$, where the base point $x_0 \in X$ is identified with $0 \in I$, the base point $y_0 \in Y$ is identified with $1 \in I$, and similarly with the other union. Since the spaces are well pointed, the quotient maps $q : X \cup I \cup Y \rightarrow X \vee Y$ and $q' : X' \cup I \cup Y' \rightarrow X' \vee Y'$ that collapse I to the common base point are homotopy equivalences. The pairs $(X \cup [0, 1], (0, 1] \cup Y)$ and $(X' \cup [0, 1], (0, 1] \cup Y')$ are excisive in $X \cup I \cup Y$ and $X' \cup I \cup Y'$, respectively. Hence, by [9, 16.24], the map $\varphi \cup \text{id}_I \cup \psi : X \cup I \cup Y \rightarrow X' \cup I \cup Y'$ is a weak homotopy equivalence. The following is clearly a commutative diagram

$$\begin{array}{ccc} X' \cup I \cup Y' & \xrightarrow{\varphi \cup \text{id}_I \cup \psi} & X \cup I \cup Y \\ q' \downarrow \simeq & & \simeq \downarrow q \\ X' \vee Y' & \xrightarrow{\varphi \vee \psi} & X \vee Y. \end{array}$$

Therefore, $\varphi \vee \psi$ is a weak homotopy equivalence. □

As a consequence, we obtain the following result.

Lemma 6.12. *Let X_α and X'_α , $\alpha \in \Lambda$ be an arbitrary family of well-pointed G -spaces, and let $\varphi_\alpha : X'_\alpha \rightarrow X_\alpha$ be an equivariant weak homotopy equivalences (mapping base points to base points). Then $\bigvee_{\alpha \in \Lambda} \varphi_\alpha : \bigvee_{\alpha \in \Lambda} X'_\alpha \rightarrow \bigvee_{\alpha \in \Lambda} X_\alpha$ is an equivariant weak homotopy equivalence.*

Proof. First notice that since the base points are fixed under the G -action, $(\bigvee_{\alpha \in \Lambda} X_\alpha)^H = \bigvee_{\alpha \in \Lambda} X_\alpha^H$, and similarly $(\bigvee_{\alpha \in \Lambda} X'_\alpha)^H = \bigvee_{\alpha \in \Lambda} X'^H_\alpha$.

Let $p(\Lambda)$ be the set of finite subsets of Λ . Then, by 6.11, for every $A \in p(\Lambda)$, $\bigvee_{\alpha \in A} \varphi^H_\alpha : \bigvee_{\alpha \in A} X'^H_\alpha \rightarrow \bigvee_{\alpha \in A} X^H_\alpha$ is a weak homotopy equivalence and induces isomorphisms between all the homotopy groups. Thus again, as in the proof of 6.10 and using [9, 15.9], $\bigvee_{\alpha \in \Lambda} \varphi^H_\alpha : \bigvee_{\alpha \in \Lambda} X'^H_\alpha \rightarrow \bigvee_{\alpha \in \Lambda} X^H_\alpha$ induces isomorphisms in all homotopy groups and hence it is a weak homotopy equivalence, thus $\bigvee_{\alpha \in \Lambda} \varphi_\alpha : \bigvee_{\alpha \in \Lambda} X'_\alpha \rightarrow \bigvee_{\alpha \in \Lambda} X_\alpha$ is an equivariant weak homotopy equivalence. □

Now we can prove the general wedge axiom for the Bredon-Illman G -equivariant homology.

Theorem 6.13. *Let X_α , $\alpha \in \Lambda$, be an arbitrary family of well-pointed G -spaces. Then there is an isomorphism $\bigoplus_{\alpha \in \Lambda} \tilde{H}^G_q(X_\alpha; M) \cong \tilde{H}^G_q(\bigvee_{\alpha \in \Lambda} X_\alpha; M)$ induced by the canonical inclusions $X_\alpha \hookrightarrow \bigvee X_\alpha$.*

Proof. For each $\alpha \in \Lambda$ there is an equivariant weak homotopy equivalence $\varphi_\alpha : \tilde{X}_\alpha \rightarrow X_\alpha$, where \tilde{X}_α is a (pointed) G -CW-complex (for instance $\tilde{X}_\alpha = |\mathcal{S}(X_\alpha)|$).

By 6.12, $\varphi = \bigvee_{\alpha} \varphi_\alpha : \bigvee_{\alpha} \tilde{X}_\alpha \rightarrow \bigvee_{\alpha} X_\alpha$ is an equivariant weak homotopy equivalence. Hence, by [5, 1.19], φ induces an isomorphism $\varphi_* : H^G_q(\bigvee_{\alpha} \tilde{X}_\alpha; M) \rightarrow H^G_q(\bigvee_{\alpha} X_\alpha; M)$. Recall that

$$H^G_q(\bigvee_{\alpha} \tilde{X}_\alpha; M) = \tilde{H}^G_q((\bigvee_{\alpha} \tilde{X}_\alpha)^+; M) \quad \text{and} \quad H^G_q(\bigvee_{\alpha} X_\alpha; M) = \tilde{H}^G_q((\bigvee_{\alpha} X_\alpha)^+; M).$$

On the other hand, the cofiber sequences

$$\mathbb{S}^0 \hookrightarrow (\bigvee_{\alpha} \tilde{X}_{\alpha})^+ \twoheadrightarrow \bigvee_{\alpha} \tilde{X}_{\alpha} \quad \text{and} \quad \mathbb{S}^0 \hookrightarrow (\bigvee_{\alpha} X_{\alpha})^+ \twoheadrightarrow \bigvee_{\alpha} X_{\alpha}$$

induce short exact sequences that fit into the diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & \tilde{H}_0^G(\mathbb{S}^0; M) & \longrightarrow & \tilde{H}_0^G((\bigvee_{\alpha} \tilde{X}_{\alpha})^+; M) & \longrightarrow & \tilde{H}_0^G(\bigvee_{\alpha} \tilde{X}_{\alpha}; M) \longrightarrow 0 \\ & & \parallel & & \cong \downarrow \varphi_*^{\dagger} & & \downarrow \varphi_* \\ 0 & \longrightarrow & \tilde{H}_0^G(\mathbb{S}^0; M) & \longrightarrow & \tilde{H}_0^G((\bigvee_{\alpha} X_{\alpha})^+; M) & \longrightarrow & \tilde{H}_0^G(\bigvee_{\alpha} X_{\alpha}; M) \longrightarrow 0 \end{array}$$

Thus, by the five-lemma, we have an isomorphism

$$\varphi_* : \tilde{H}_0^G(\bigvee_{\alpha} \tilde{X}_{\alpha}; M) \longrightarrow \tilde{H}_0^G(\bigvee_{\alpha} X_{\alpha}; M).$$

On the other hand, since $\tilde{H}_q^G(\mathbb{S}^0; M) = 0$ if $q > 0$, there are isomorphisms

$$\tilde{H}_q^G((\bigvee_{\alpha} \tilde{X}_{\alpha})^+; M) \cong \tilde{H}_q^G(\bigvee_{\alpha} \tilde{X}_{\alpha}; M)$$

$$\tilde{H}_q^G((\bigvee_{\alpha} X_{\alpha})^+; M) \cong \tilde{H}_q^G(\bigvee_{\alpha} X_{\alpha}; M)$$

and thus also an isomorphism

$$\varphi_* : \tilde{H}_q^G(\bigvee_{\alpha} \tilde{X}_{\alpha}; M) \longrightarrow \tilde{H}_q^G(\bigvee_{\alpha} X_{\alpha}; M).$$

Similarly, for every $\alpha \in \Lambda$, there are isomorphisms

$$\varphi_{\alpha*} : \tilde{H}_q^G(\tilde{X}_{\alpha}; M) \longrightarrow \tilde{H}_q^G(X_{\alpha}; M).$$

Since the G -spaces \tilde{X}_{α} are G -CW-complexes, by 6.10 there is an isomorphism

$$\bigoplus_{\alpha \in \Lambda} \tilde{H}_q^G(\tilde{X}_{\alpha}; M) \cong \tilde{H}_q^G(\bigvee_{\alpha} \tilde{X}_{\alpha}; M)$$

for all $q \geq 0$, and by the isomorphisms above it follows that there is an isomorphism

$$\bigoplus_{\alpha \in \Lambda} \tilde{H}_q^G(X_{\alpha}; M) \cong \tilde{H}_q^G(\bigvee_{\alpha} X_{\alpha}; M)$$

as desired. □

Next we make some calculations using the homotopical approach to Bredon-Illman homology.

Proposition 6.14. *Let X be a pointed G -space of the homotopy type of a G -CW-complex. Assume that X^H is connected for each $H \subset G$. Then*

$$\tilde{H}_0^G(X; M) = 0.$$

Proof. By Lemma 6.1, $C(X^H) = \pi_0(X^H)$ for each $H \subset G$. Hence

$$\pi_0(F(X^H, M(G/H))) \cong \tilde{H}_0(X^H; M(G/H)) = 0.$$

Thus the topological groups $F(X^H, M(G/H))$ are path-connected and so is their product. Since $\mathcal{F}^G(X, M)$ is a quotient of this product, it is also path-connected. Hence $0 = \pi_0(\mathcal{F}^G(X, M)) \cong \tilde{H}_0^G(X; M)$. \square

Proposition 6.15. *Let X be a G -space of the homotopy type of a G -CW-complex. Assume that X^H is connected for each $H \subset G$. Consider the family \mathcal{H} of all $H \subset G$ such that $X^H \neq \emptyset$. Then $H_0^G(X; M)$ is a quotient of $\bigoplus_{H \in \mathcal{H}} M(G/H)$. Furthermore, if X has a fixed point, then $H_0^G(X; M) \cong M(G/G)$.*

Proof. For each $H \in \mathcal{H}$, there is an isomorphism

$$\pi_0(F((X^H)^+, M(G/H))) \cong H_0(X^H; M(G/H)) \cong M(G/H).$$

Hence one has an epimorphism

$$\bigoplus_{H \in \mathcal{H}} M(G/H) \cong \pi_0\left(\prod_{H \in \mathcal{H}} F((X^H)^+, M(G/H))\right) \rightarrow \pi_0(\mathcal{F}^G(X^+, M)) \cong H_0^G(X; M).$$

Furthermore, if X has a fixed point x_0 , then $X^+ = X \vee \mathbb{S}^0$, where we take x_0 as base point of X . By the exactness axiom, one always has $\tilde{H}_*^G(X \vee Y; M) \cong \tilde{H}_*^G(X; M) \oplus \tilde{H}_*^G(Y; M)$. Then, in particular, $\tilde{H}_0^G(X^+; M) \cong \tilde{H}_0^G(X; M) \oplus \tilde{H}_0^G(\mathbb{S}^0; M)$. Since by the previous proposition $\tilde{H}_0^G(X; M) = 0$, one obtains

$$H_0^G(X; M) \cong \tilde{H}_0^G(\mathbb{S}^0; M) \cong M(G/G).$$

\square

7. Dold-Thom topological groups and the transfer

In this section we summarize the properties of the topological groups defined and explain which of them admit a transfer, either for ordinary or for ramified covering G -maps. We start with the following concept.

Definition 7.1. Let M be a covariant coefficient system for G . We shall call a functor $\mathcal{G}(-, M) : G\text{-Top} \rightarrow \text{Topab}$ a *Dold-Thom topological group functor with coefficients in M for a subcategory $\mathcal{T} \subset G\text{-Top}$* if there is a natural isomorphism

$$\varphi_X : \pi_*(\mathcal{G}(X, M)) \rightarrow \tilde{H}_*^G(X; M),$$

where the right-hand side denotes the reduced G -equivariant Bredon-Illman homology groups of X with coefficients in M , and X is an object of \mathcal{T} .

EXAMPLES 7.2. The following are examples of Dold-Thom topological group functors with coefficients in a coefficient system M :

1. The groups $F^G(X, M) = |F^G(\mathcal{S}(X), M)|$, for any G -space X , defined in [2].
2. The groups $\mathbb{F}^G(X, M)$ defined in Section 5 of [3], when M is a homological Mackey functor and X has the homotopy type of a G -CW-complex.

3. The groups $\mathcal{F}^G(X, M)$ as shown in 3.11.
4. The groups $|F^G(Q, M)|$ for $X = |Q|$ and Q a simplicial G -set, in particular, the simplicial G -set associated to a simplicial G -complex as shown in [5].
5. The groups $\text{AG}(X)$ and $\text{AG}(X; m)$ defined by Dold and Thom [7] in the nonequivariant case, where the coefficients are a cyclic group (\mathbb{Z} and $\mathbb{Z}/m\mathbb{Z}$, respectively) and X is a (countable) simplicial complex. Here G does not stand for any group.
6. The groups $B(L, X)$ defined by McCord [15] in the nonequivariant case and coefficients in an abelian group L , where X is a weak Hausdorff k -space of the homotopy type of a CW-complex.
7. The groups $\text{AG}(X)$ defined by Lima-Filho [12] for a G -CW-complex X and coefficients in \mathbb{Z} .
8. The groups $L \otimes X$ defined by dos Santos [10] for a pointed G -CW-complex X and coefficients in a G -module L .
9. The groups $\mathcal{G}X \otimes_{G\mathcal{F}} M$ defined in [17] for a pointed G -CW-complex X .

And if M takes values in $k\text{-Mod}$, where k is a field of characteristic 0 or a prime p that does not divide $|G|$:

10. The groups $\overline{F}^G(X, M)$, by 5.2(a) and 1.
11. The groups $\overline{\mathbb{F}}^G(X, M)$, if X has the homotopy type of a G -CW-complex, by 5.3 and 2.
12. The groups $\overline{\mathcal{F}}^G(X, M)$ for a strong ρ -space X , by 5.2(b), 5.3, 4.6, and 3.

Definition 7.3. Let M be a Mackey functor for the finite group G and $\mathcal{G}(-, M)$ be a Dold-Thom topological group functor with coefficients in (the covariant part of) M . Let $p : E \rightarrow X$ be either an n -fold G -equivariant ordinary covering map (see [2]) or an n -fold G -equivariant ramified covering map (see [4]). By a *transfer* for p in $\mathcal{G}(-, M)$ we understand a continuous homomorphism

$$t_p^G : \mathcal{G}(X^+, M) \rightarrow \mathcal{G}(E^+, M),$$

which satisfies the following conditions:

- (a) Pullback: If $f : Y \rightarrow X$ is continuous and we take the pullback diagram

$$\begin{array}{ccc} E' & \xrightarrow{\tilde{f}} & E \\ q \downarrow & & \downarrow p \\ Y & \xrightarrow{f} & X, \end{array}$$

then $t_p^G \circ f_*^G = \tilde{f}_*^G \circ t_q^G : \mathcal{G}(Y^+, M) \rightarrow \mathcal{G}(E^+, M)$.

- (b) Normalization: If $p = \text{id}_X : X \rightarrow X$, then $t_{\text{id}_X}^G = \text{id} : \mathcal{G}(X^+, M) \rightarrow \mathcal{G}(X^+, M)$.

- (c) Functoriality: If $p : E \rightarrow X$ and $q : X \rightarrow Y$ are G -equivariant ordinary, resp. ramified covering maps, then

$$t_{q \circ p}^G = t_p^G \circ t_q^G : \mathcal{G}(Y^+, M) \rightarrow \mathcal{G}(E^+, M).$$

- (d) If M is homological, then the composite $p_*^G \circ t_p^G : \mathcal{G}(X^+, M) \rightarrow \mathcal{G}(X^+, M)$ is multiplication by n .

EXAMPLES 7.4. The following Dold-Thom topological group functors have transfers for p :

1. If $p : E \rightarrow X$ is an n -fold G -equivariant ordinary covering map and M is any Mackey functor, then there is a transfer $t_p^G : F^G(X^+, M) \rightarrow F^G(E^+, M)$, as shown in [2].
2. If $p : E \rightarrow X$ is an n -fold G -equivariant ramified covering map between strong ρ -spaces of the homotopy type of G -CW-complexes, and M is homological, then there is a transfer $t_p^G : \mathbb{F}^G(X^+, M) \rightarrow \mathbb{F}^G(E^+, M)$, as shown in [4].
3. If $p : K \rightarrow Q$ is a special G -equivariant simplicial ramified covering map and M is any Mackey functor, then there is a transfer $|t_p^G| : |F^G(Q^+, M)| \rightarrow |F^G(K^+, M)|$, as shown in [5].
4. If $p : E \rightarrow X$ is an n -fold G -equivariant ramified covering map between strong ρ -spaces of the homotopy type of G -CW-complexes, and M is homological, then there is a transfer $t_p^G : \mathcal{F}^G(X^+, M) \rightarrow \mathcal{F}^G(E^+, M)$, by 4.3 and Example 2.

REMARK 7.5. Let $p : E \rightarrow X$ be an n -fold G -equivariant covering map. The restrictions $p^H : E^H \rightarrow X^H$ are not, in general, covering maps. Thus there are no transfers $F(X^{H+}, M(G/H)) \rightarrow F(E^{H+}, M(G/H))$. Since the topology of the groups $\mathcal{F}^G(X^+, M)$ and $\mathcal{F}^G(E^+, M)$ is given in terms of that of the groups $F(X^{H+}, M(G/H))$ and $F(E^{H+}, M(G/H))$, it does not seem possible to prove the continuity of t_p^G . And even if the transfers t_{p^H} exist, they do not commute with the identifications. However, if as stated in Example 4 above, the spaces are ρ -spaces, then the groups $\mathcal{F}^G(X^+, M)$ and $\mathcal{F}^G(E^+, M)$ coincide with the groups $\mathbb{F}^G(X^+, M)$ and $\mathbb{F}^G(E^+, M)$, as shown in 4.4, and one can show that the transfer is continuous.

Now we study the homotopy type of the Dold-Thom topological groups. First we have the following general result.

Theorem 7.6. *Let A be a locally connected topological abelian group in the category $k\text{-Top}$ that has the homotopy type of a CW-complex. Then there is a homotopy equivalence*

$$A \simeq \bigoplus_{q \geq 0} K(\pi_q(A), q),$$

where $K(\pi_q(A), q)$ denotes the corresponding Eilenberg-Mac Lane space.

Proof. Since translation by a_0 , $a_0 \in A$, is a homeomorphism in the k -topology, the connected component A_0 of $0 \in A$ is a closed subgroup of A .

By Lemma 6.1, the connected components of A coincide with the path-components. Thus A is the topological sum of all its path-components, and they are closed and open, because A is locally connected. Since all path-components of A are homeomorphic (via translation) to A_0 , we have a homeomorphism

$$A \cong \pi_0(A) \times A_0. \tag{7.7}$$

Since A has the homotopy type of a CW-complex, so does A_0 . Consider

$$A_0 \xrightarrow{i} F(A_0, \mathbb{Z}) \xrightarrow{\nu} A_0,$$

given by $i(a) = 1a$ and $\nu(u) = \sum_{a \in A_0} u(a)a$. Then i and ν are clearly continuous, and the composite $\nu \circ i$ is equal to the identity id_{A_0} . Applying the functor π_q we have the following

$$\begin{array}{ccccc} & & 1 & & \\ & \curvearrowright & & \curvearrowleft & \\ \pi_q(A_0) & \xrightarrow{i\#} & \pi_q(F(A_0, \mathbb{Z})) & \xrightarrow{\nu\#} & \pi_q(A_0) \\ & \searrow h_q & \downarrow \cong & \swarrow \lambda_q & \\ & & \tilde{H}_q(A_0; \mathbb{Z}) & & \end{array}$$

where λ_q is defined so that the triangle commutes, and thus we obtain a left inverse for the Hurewicz homomorphism.

Now consider a homotopy equivalence $A_0 \xrightarrow{\varphi} C_0$, where C_0 is a CW-complex, and the diagram

$$\begin{array}{ccccc} \pi_q(A_0) & \xrightarrow{h_q} & \tilde{H}_q(A_0) & \xrightarrow{\lambda_q} & \pi_q(A_0) \\ \varphi\# \downarrow \cong & & \varphi\# \downarrow \cong & & \varphi\# \downarrow \cong \\ \pi_q(C_0) & \xrightarrow{h_q} & \tilde{H}_q(C_0) & \xrightarrow{\alpha_q} & \pi_q(C_0), \end{array}$$

where α_q is such that the diagram commutes. Hence α_q is a left inverse to Hurewicz too. Since C_0 is a connected CW-complex, by Moore's theorem (see [19, IX(1.9)]), we have $C_0 \simeq \bigoplus_{q \geq 1} K(\pi_q(C_0), q)$ (see 6.2). Hence, using (7.7), we get

$$A \cong \pi_0(A) \times A_0 \simeq \pi_0(A) \times \bigoplus_{q \geq 1} K(\pi_q(C_0), q) \approx \bigoplus_{q \geq 0} K(\pi_q(A_0), q).$$

□

REMARK 7.8. This result was proved by John Moore in [16] for the case of connected simplicial abelian groups and by Dold and Thom [8] for the case of connected finite polyhedral abelian groups.

As a consequence of Theorem 7.6 we have the following.

Theorem 7.9. *Let $\mathcal{G}(-, M)$ is a Dold-Thom topological group functor. If X is a pointed G -space such that $\mathcal{G}(X, M)$ is locally connected and has the homotopy type*

of a CW-complex, then

$$\mathfrak{S}(X, M) \simeq \bigoplus_{q \geq 0} K(\tilde{H}_q^G(X; M), q).$$

Hence these groups are unique up to homotopy. \square

Proposition 7.10. (a) *The topological groups $F^G(X, M)$ are CW-complexes for any pointed G -space X .*

- (b) *If X is a pointed G -CW-complex, then $\mathcal{F}^G(X, M)$ is locally connected and has the homotopy type of a CW-complex.*
- (c) *If X is a pointed G -simplicial complex or a finite-dimensional countable locally finite G -CW-complex, then $\mathbb{F}^G(X, M)$ is locally connected and has the homotopy type of a CW-complex.*

We conclude that in all these cases the topological groups have the homotopy type of $\bigoplus_{q \geq 0} K(\tilde{H}_q^G(X; M), q)$.

Proof. (a) follows from the fact that $F^G(X, M)$ is the geometric realization $|F^G(\mathfrak{S}(X), M)|$.

(b) First notice that the property of a space being locally connected is inherited by quotient spaces. Hence, if X is locally connected, so is also $\prod_{n \geq 1} (L \times X)^n$ for any abelian group L , and given the quotient map $\prod_{n \geq 1} (L \times X)^n \twoheadrightarrow F(X, L)$, the topological group $F(X, L)$ is locally connected.

Now, if X is a G -CW-complex, then X^H is locally connected for every $H \subset G$. Hence each topological group $F(X^H, M(G/H))$ is locally connected and since by definition there is a quotient map $\prod_{H \subset G} F(X^H, M(G/H)) \twoheadrightarrow \mathcal{F}^G(X, M)$, the topological group $\mathcal{F}^G(X, M)$ is also locally connected. Furthermore there is a G -homotopy equivalence $\rho_X : |\mathfrak{S}(X)| \longrightarrow X$, which by the homotopy invariance 3.4 induces a homotopy equivalence $\rho_{X*}^G : \mathcal{F}^G(|\mathfrak{S}(X)|, M) \longrightarrow \mathcal{F}^G(X, M)$. But by 3.10, there is an isomorphism of topological groups $\mathcal{F}^G(|\mathfrak{S}(X)|, M) \cong |F^G(\mathfrak{S}(X), M)|$, thus the first is a CW-complex, and hence $\mathcal{F}^G(X, M)$ has the homotopy type of a CW-complex.

(c) Actually we only need X to be a pointed G -CW-complex which is also a strong ρ -space, then by 4.4, $\mathbb{F}^G(X, M) = \mathcal{F}^G(X, M)$ and the result follows from (b).

In any case, the corresponding topological group satisfies the assumptions of Theorem 7.6 and we obtain the conclusion. \square

References

- [1] M. Aguilar and C. Prieto, *Topological abelian groups and equivariant homology*, Communications in Algebra **36** (2008), 434–454.
- [2] M. Aguilar and C. Prieto, *Equivariant homotopical homology with coefficients in a Mackey functor*, Topology and its Applications **154** (2007), 2826–2848 (with a correction, *ibid.* **156** (2009), 1609–1613).
- [3] M. Aguilar and C. Prieto, *Topological groups and Mackey functors*, Bol. Soc. Mat. Mex. **14** (2008), 233–262.
- [4] M. Aguilar and C. Prieto, *The transfer for ramified covering G -maps*, Forum Math. **22** (2010) 1089–1115.
- [5] M. Aguilar and C. Prieto, *Bredon homology and ramified covering G -maps*, Topology and its Applications **157** (2010) 401–416.
- [6] T. tom Dieck, *Transformation Groups*, Walter de Gruyter, Berlin & New York, 1987.
- [7] A. Dold and R. Thom, *Quasifaserungen und unendliche symmetrische Produkte*, Annals of Math. **67** (1958), 239–281.
- [8] A. Dold and R. Thom, *Une généralisation de la notion d'espace fibré. Application aux produits symétriques infinis*, C. R. Acad. Sci. Paris **242** (1956), 1680–1682.
- [9] B. Gray, *Homotopy Theory*, Academic Press New York San Francisco London 1975.
- [10] P. F. dos Santos, *A note on the equivariant Dold-Thom theorem*, J. Pure Appl. Algebra **183** (2003), 299–312.
- [11] S. Illman, *Equivariant singular homology and cohomology I*, Memoirs of the Amer. Math. Soc. Vol. 1, Issue 2, No. 156, 1975.
- [12] P. Lima-Filho, *On the equivariant homotopy of free abelian groups on G -spaces and G -spectra*, Math. Z. **224** (1997), 567–601.
- [13] J. P. May, *Simplicial Objects in Algebraic Topology*, The University of Chicago Press, Chicago London 1992.
- [14] J. P. May, *A Concise Course in Algebraic Topology*, The University of Chicago Press, Chicago & London 1999.
- [15] M. C. McCord, *Classifying spaces and infinite symmetric products*, Trans. Amer. Math. Soc. **146** (1969), 273–298.
- [16] J. C. Moore, *Semi-simplicial complexes and Postnikov systems*, Symposium Intern. de Topología Algebraica, 232–247, UNAM y UNESCO, México 1958.
- [17] Z. Nie, *A functor converting equivariant homology to homotopy*, Bull. London Math. Soc., **39**(3) (2007), 499–508.

- [18] R. M. Vogt, *Convenient categories of topological spaces for homotopy theory*, Archiv der Mathematik **XXII** (1971), 545–555.
- [19] G. W. Whitehead, *Elements of Homotopy Theory*, Springer-Verlag, New York Heidelberg Berlin 1978.

<http://www.emis.de/ZMATH/>
<http://www.ams.org/mathscinet>

This article may be accessed via WWW at <http://tcms.org.ge/Journals/JHRS/>

Marcelo A. Aguilar
marcelo@math.unam.mx
Instituto de Matemáticas
Universidad Nacional Autónoma de México
04510 México, D.F., Mexico

Carlos Prieto
cprieto@math.unam.mx
Instituto de Matemáticas
Universidad Nacional Autónoma de México
04510 México, D.F., Mexico