DE GRUYTER
OPEN

DOI 10.2478/tmj-2018-0008

Existence of solution for a coupled system of Urysohn-Stieltjes functional integral equations

A. M. A. El-Sayed[1] and M. M. A. Al-Fadel[2]

[1]Faculty of Science, Alexandria University, Alexandria, Egypt
[2]Faculty of Science, Omar Al-Mukhtar University, Libya
E-mail: amasayed@alexu.edu.eg[1], najemeoe1234@gmail.com[2]

Abstract

We present an existence theorem for at least one continuous solution for a coupled system of nonlinear functional (delay) integral equations of Urysohn-Stieltjes type.

2010 Mathematics Subject Classification. **74H10**. 45G10, 47H30
Keywords. Coupled system, Urysohn-Stieltjes integral equations, continuous solution.

1 Introduction and preliminaries

Urysohn-Stieltjes integral operators and Urysohn-Stieltjes integral equations have been studied by some authors (see [1]-[8]). The coupled system of integral equations have been studied, recently, by some authors (see [10]-[12]). Our aim here is to study the existence of at least one solution for a coupled system of nonlinear functional (delay) integral equations of Urysohn-Stieltjes type in the space of continuous functions.

In what follows let $I = [0,1]$ be a fixed interval. Denote by $C(I) = C[0,1]$ the Banach space consisting of all continuous functions acting from the interval I into R with the standard norm

$$\| x \| = \sup_{t \in I} | x(t) | .$$

Consider the nonlinear Urysohn-Stieltjes integral equation

$$x(t) = p(t) + \int_0^1 f(t,s,x(s)) \; d_s g(t,s), \; t \in I = [0,1] \tag{1}$$

where $g : I \times I \to R$ and the symbol d_s indicates the integration with respect to s.
Equations of type (1) and some of their generalizations were considered in paper (see [3]), for the properties of the Urysohn-Stieltjes integral (see Banaś [1]).

In this paper, we generalize this result for the coupled system of Urysohn-Stieltjes functional (delay) integral equations

$$\begin{aligned} x(t) &= p_1(t) + \int_0^1 f_1(t,s,y(\psi_1(s))) \; d_s g_1(t,s), \; t \in I \\ y(t) &= p_2(t) + \int_0^1 f_2(t,s,x(\psi_2(s))) \; d_s g_2(t,s), \; t \in I \end{aligned} \tag{2}$$

Tbilisi Mathematical Journal 11(1) (2018), pp. 117–125.
Tbilisi Centre for Mathematical Sciences.
Received by the editors: 14 August 2017.
Accepted for publication: 03 January 2018.

Unauthenticated
Download Date | 3/3/18 2:38 PM

in the Banach space $C(I)$.

2 Existence of solutions

In this section we discuss the existence of solutions for the coupled system of nonlinear Urysohn-Stieltjes integral equations in C(I). In our further considerations, we shall assume that the following conditions are satisfied:

(i) $p_i : I \to R$ are continuous functions on I, $p = \sup_t \mid p_i(t) \mid$, $i = 1, 2$.

(ii) $\psi_i : I \to I$ are continuous functions such that $\psi_i(t) \leq t$, $i = 1, 2$.

(iii) $f_i : I \times I \times R \to R$, $i = 1, 2$ are continuous functions such that there exist continuous functions $a_i : I \times I \to I$ and continuous and nondecreasing functions $\varphi_i : R_+ \to R_+$ such that

$$\mid f_i(t, s, x) \mid \leq a_i(t, s)\varphi_i(\mid x \mid)$$

for $t, s \in I$, $x \in R$, $i = 1, 2$. Moreover, we put $k = \max\{a_i(t, s) : t, s \in I, i = 1, 2\}$.

(iv) $g_i : I \times I \to R$, $i = 1, 2$ and for all $t_1, t_2 \in I$ with $t_1 < t_2$, and the function $s \to g_i(t_2, s) - g_i(t_1, s)$ is nondecreasing on I.

(v) $g_i(0, s) = 0$ for any $s \in I$.

(vi) The functions $t \to g_i(t, 1)$ and $t \to g_i(t, 0)$ are continuous on I. Put

$$\mu = \max\{\sup \mid g_i(t, 1) \mid + \sup \mid g_i(t, 0) \mid \ on\ I\}, \quad i = 1, 2.$$

(vii) There exists a positive number r satisfying the inequality

$$p + (k\varphi_i(r))\mu \leq r.$$

Remark 2.1. Observe that Assumptions (iv) and (v) imply that the function $s \to g(t, s)$ is nondecreasing on the interval I, for any fixed $t \in I$ (Remark 1 in [4]). Indeed, putting $t_2 = t$, $t_1 = 0$ in (iv) and keeping in mind (v), we obtain the desired conclusion. From this observation, it follows immediately that, for every $t \in I$, the function $s \to g(t, s)$ is of bounded variation on I.

Now, let X be the Banach space of all ordered pairs (x, y), $x, y \in C(I)$ with the norm

$$\| (x, y) \|_X = \max\{||x||, ||y||\}$$

where

$$\| x \| = \sup_{t \in I} \mid x(t) \mid, \qquad \| y \| = \sup_{t \in I} \mid y(t) \mid .$$

It is clear that $(X, ||.||_X)$ is Banach space.

Theorem 2.2. Let the assumptions (i)-(vii) be satisfied, then the coupled system (2) has at least one solution in X.

Unauthenticated
Download Date | 3/3/18 2:38 PM

Proof. Define the operator T by

$$T(x,y)(t) = (T_1y(t), T_2x(t))$$

where

$$T_1y(t) = x(t) = p_1(t) + \int_0^1 f_1(t,s,y(\psi_1(s)))\ d_sg_1(t,s)$$

$$T_2x(t) = y(t) = p_2(t) + \int_0^1 f_2(t,s,x(\psi_2(s)))\ d_sg_2(t,s).$$

We prove a few results concerning the continuity and compactness of these operators in the space of continuous functions.
We define the set U by

$$U = \{u = (x(t), y(t)) \mid (x(t), y(t)) \in X :\| (x,y) \|_X \leq r\}$$

Let $(x,y) \in U$ and define

$$\theta(\varepsilon) = \sup\{\mid f_1(t_2,s,y) - f_1(t_1,s,y) \mid, \mid f_2(t_2,s,x) - f_2(t_1,s,x) \mid : t_1,t_2 \in I,\ \ \mid t_2 - t_1 \mid \leq \varepsilon,\ x \in R\}.$$

Now, for $(x,y) \in U$, we have

$$\begin{aligned}
\mid T_1y(t) \mid \quad &\leq \quad \mid p_1(t) \mid + \mid \int_0^1 f_1(t,s,y(\psi_1(s)))\ d_sg_1(t,s) \mid \\
&\leq \quad \sup_t \mid p_1(t) \mid + \int_0^1 \mid f_1(t,s,y(\psi_1(s))) \mid\ \ d_s(\bigvee_{z=0}^{s} g_1(t,z)) \\
&\leq \quad p + \int_0^1 (a_1(t,s)\varphi_1(\mid y(\psi_1(s)) \mid))\ d_s(\bigvee_{z=0}^{s} g_1(t,z)) \\
&\leq \quad p + (k\varphi_1(\sup_s \mid y(\psi_1(s)) \mid)) \int_0^1 d_sg_1(t,s) \\
&\leq \quad p + (k\varphi_1(\| y \|))[g_1(t,1) - g_1(t,0)] \\
&\leq \quad p + (k\varphi_1(r))[\mid g_1(t,1) \mid + \mid g_1(t,0) \mid] \\
&\leq \quad p + (k\varphi_1(r))[\sup_t \mid g_1(t,1) \mid + \sup_t \mid g_1(t,0) \mid] \\
&\leq \quad p + (k\varphi_1(r))\mu
\end{aligned}$$

then

$$\| T_1y \| \ \leq\ p + (k\varphi_1(r))\mu.$$

By a similar way can deduce that

$$\| T_2x \| \ \leq\ p + (k\varphi_2(r))\mu.$$

Therefore,

$$\| Tu \|_X = \| T(x,y) \|_X = \| (T_1y, T_2x) \|_X = \max\{\| T_1y \|, \| T_2x \|\} \leq r.$$

Unauthenticated
Download Date | 3/3/18 2:38 PM

Thus for every $u=(x,y)\in U$, we have $Tu\in U$ and hence $TU\subset U$, (i.e $T:U\to U$).
This means that the functions of TU are uniformly bounded on I, it is clear that the set U is nonempty, bounded, closed and convex.

Now, we prove that the set TU is relatively compact.
For $u=(x,y)\in U$, for all $\varepsilon>0$, $\delta>0$ and for each $t_1,t_2\in I$, and $t_1<t_2$ such that $\mid t_2-t_1\mid<\delta$, then

$$\begin{aligned}
\mid T_1y(t_2)-T_1y(t_1)\mid &\leq \mid p_1(t_2)-p_1(t_1)\mid \\
&+ \mid\int_0^1 f_1(t_2,s,y(\psi_1(s)))\ d_sg_1(t_2,s)-\int_0^1 f_1(t_1,s,y(\psi_1(s)))\ d_sg_1(t_1,s)\mid \\
&\leq \mid p_1(t_2)-p_1(t_1)\mid \\
&+ \mid\int_0^1 f_1(t_2,s,y(\psi_1(s)))\ d_sg_1(t_2,s)-\int_0^1 f_1(t_1,s,y(\psi_1(s)))\ d_sg_1(t_2,s)\mid \\
&+ \mid\int_0^1 f_1(t_1,s,y(\psi_1(s)))\ d_sg_1(t_2,s)-\int_0^1 f_1(t_1,s,y(\psi_1(s)))\ d_sg_1(t_1,s)\mid \\
&\leq \mid p_1(t_2)-p_1(t_1)\mid \\
&+ \mid\int_0^1 [f_1(t_2,s,y(\psi_1(s)))-f_1(t_1,s,y(\psi_1(s)))]\ d_sg_1(t_2,s)\mid \\
&+ \mid\int_0^1 f_1(t_1,s,y(\psi_1(s)))\ d_s(g_1(t_2,s)-g_1(t_1,s))\mid \\
&\leq \mid p_1(t_2)-p_1(t_1)\mid \\
&+ \int_0^1 \mid f_1(t_2,s,y(\psi_1(s)))-f_1(t_1,s,y(\psi_1(s)))\mid\ d_s(\bigvee_{z=0}^{s} g_1(t_2,z)) \\
&+ \int_0^1 \mid f_1(t_1,s,y(\psi_1(s)))\mid\ d_s(\bigvee_{z=0}^{s}[g_1(t_2,z)-g_1(t_1,z)]) \\
&\leq \mid p_1(t_2)-p_1(t_1)\mid +\int_0^1 \theta(\varepsilon)\ d_s(\bigvee_{z=0}^{s} g_1(t_2,z)) \\
&+ \int_0^1 (a_1(t_1,s)\varphi_1(\mid y(\psi_1(s))\mid))\ d_s(\bigvee_{z=0}^{s}[g_1(t_2,z)-g_1(t_1,z)]) \\
&\leq \parallel p_1(t_2)-p_1(t_1)\parallel +\theta(\varepsilon)\int_0^1 d_s(g_1(t_2,s)) \\
&+ (k\varphi_1(\parallel y\parallel))\int_0^1 d_s[g_1(t_2,s)-g_1(t_1,s)]
\end{aligned}$$

Unauthenticated
Download Date | 3/3/18 2:38 PM

$$\begin{aligned} &\leq \| p_1(t_2) - p_1(t_1) \| + \theta(\varepsilon)[g_1(t_2,1) - g_1(t_2,0)] \\ &+ (k\varphi_1(r))\{[g_1(t_2,1) - g_1(t_1,1)] - [g_1(t_2,0) - g_1(t_1,0)]\} \\ &\leq \| p_1(t_2) - p_1(t_1) \| + \theta(\varepsilon)[g_1(1,1) - g_1(1,0)] \\ &+ (k\varphi_1(r))\{[\mid g_1(t_2,1) - g_1(t_1,1) \mid + \mid g_1(t_2,0) - g_1(t_1,0) \mid]\} \end{aligned}$$

Hence

$$\begin{aligned} \| T_1y(t_2) - T_1y(t_1) \| &\leq \| p_1(t_2) - p_1(t_1) \| + \theta(\varepsilon)[g_1(1,1) - g_1(1,0)] \\ &+ (k\varphi_1(r))[\mid g_1(t_2,1) - g_1(t_1,1) \mid + \mid g_1(t_2,0) - g_1(t_1,0) \mid]. \end{aligned}$$

As done above we can obtain

$$\begin{aligned} \| T_2x(t_2) - T_2x(t_1) \| &\leq \| p_2(t_2) - p_2(t_1) \| + \theta(\varepsilon)[g_2(1,1) - g_2(1,0)] \\ &+ (k\varphi_2(r))[\mid g_2(t_2,1) - g_2(t_1,1) \mid + \mid g_2(t_2,0) - g_2(t_1,0) \mid]. \end{aligned}$$

Now, from the definition of the operator T we get

$$\begin{aligned} Tu(t_2) - Tu(t_1) &= T(x,y)(t_2) - T(x,y)(t_1) \\ &= (T_1y(t_2), T_2x(t_2)) - (T_1y(t_1), T_2x(t_1)) \\ &= (T_1y(t_2) - T_1y(t_1), T_2x(t_2) - T_2x(t_1)) \end{aligned}$$

Therefore,

$$\begin{aligned} \| Tu(t_2) - Tu(t_1) \|_X &= \| (T_1y(t_2) - T_1y(t_1), T_2x(t_2) - T_2x(t_1)) \|_X \\ &= \max\{\| T_1y(t_2) - T_1y(t_1) \|, \| T_2x(t_2) - T_2x(t_1) \|\} \\ &\leq \max\{\| p_1(t_2) - p_1(t_1) \| + \theta(\varepsilon)[g_1(1,1) - g_1(1,0)] \\ &+ (k + rb)[\mid g_1(t_2,1) - g_1(t_1,1) \mid + \mid g_1(t_2,0) - g_1(t_1,0) \mid] \\ &, \| p_2(t_2) - p_2(t_1) \| + \theta(\varepsilon)[g_2(1,1) - g_2(1,0)] \\ &+ (k + rb)[\mid g_2(t_2,1) - g_2(t_1,1) \mid + \mid g_2(t_2,0) - g_2(t_1,0) \mid]\}. \end{aligned}$$

This means that the class of $\{Tu(t)\}$ is equi-continuous on I, then by Arzéla-Ascoil theorem TU is relatively compact.

Now, we will show that the operator $T : U \to U$ is continuous.
Firstly, we prove that T_1 is continuous, for all $\varepsilon > 0$ and $\delta > 0$, let $y_1(t)$ and $y_2(t) \in C[0,1]$ and $\mid y_1(t) - y_2(t) \mid < \delta$, then

$$\begin{aligned} \mid T_1y_1(t) - T_1y_2(t) \mid &\leq \mid \int_0^1 f_1(t,s,y_1(\psi_1(s)))\ d_sg_1(t,s) - \int_0^1 f_1(t,s,y_2(\psi_1(s)))\ d_sg_1(t,s) \mid \\ &\leq \int_0^1 \mid f_1(t,s,y_1(\psi_1(s))) - f_1(t,s,y_2(\psi_1(s))) \mid\ d_s(\bigvee_{z=0}^{s} g_1(t,z)) \\ &\leq \varepsilon^* \int_0^1 d_s(\bigvee_{z=0}^{s} g_1(t,z)) \end{aligned}$$

Unauthenticated
Download Date | 3/3/18 2:38 PM

$$\begin{aligned} &\leq \varepsilon^* \int_0^1 d_s g_1(t,s) \\ &\leq \varepsilon^* \, [g_1(t,1) - g_1(t,0)] \\ &\leq \varepsilon^* \, [\mid g_1(t,1) \mid + \mid g_1(t,0) \mid] \\ &\leq \varepsilon^* \, \mu = \varepsilon \end{aligned}$$

Therefore,

$$\mid T_1 y_1(t) - T_1 y_2(t) \mid \leq \varepsilon.$$

This means that the operator T_1 is continuous.
By a similar way as done above we can prove that for any $x_1(t),\ x_2(t) \in C[0,1]$ and $\mid x_1(t) - x_2(t) \mid < \delta$, we have

$$\mid T_2 x_1(t) - T_2 x_2(t) \mid \leq \varepsilon.$$

Hence T_2 is continuous operator.
The operators T_1 and T_2 are continuous operators imply that T is continuous operator.
Since all conditions of Schauder fixed point theorem are satisfied, then T has at least one fixed point $u = (x, y) \in U$, which completes the proof. ■

Corollary 2.3. Under the assumptions of Theorem 2.2 (with $g_i(t,s) = g_i(s)$), the coupled system of Urysohn-Stieltjes integral equations

$$x(t) = p_1(t) + \int_0^1 f_1(t, s, y(\psi_1(s)))\ d_s g_1(s),\ t \in I$$

$$y(t) = p_2(t) + \int_0^1 f_2(t, s, x(\psi_2(s)))\ d_s g_2(s),\ t \in I$$

has a solution $u = (x, y) \in U$.

In what follows, we provide some examples illustrating the above obtained results.
Example 1. Consider the functions $g_i : I \times I \to R$ defined by the formula

$$\begin{aligned} g_1(t,s) &= \begin{cases} t \ln \frac{t+s}{t}, & \text{for } t \in [0,1], \quad s \in I, \\ 0, & \text{for } t = 0, \quad s \in I. \end{cases} \\ g_2(t,s) &= t(t+s-1), \quad t \in I. \end{aligned}$$

It can be easily seen that the functions $g_1(t,s)$ and $g_2(t,s)$ satisfies assumptions (iv)-(vi) given in Theorem 1. In this case, the coupled system of Urysohn-Stieltjes integral equations (2) has the form

$$\begin{aligned} x(t) &= p_1(t) + \int_0^1 \frac{t}{t+s} f_1(t, s, y(\psi_1(s)))\ ds, \quad t \in I \\ y(t) &= p_2(t) + \int_0^1 t f_2(t, s, x(\psi_2(s)))\ ds, \quad t \in I, \end{aligned} \tag{3}$$

Unauthenticated
Download Date | 3/3/18 2:38 PM

Therefore, the coupled system (3) has at least one solution $x, y \in C[0,1]$.

Example 2. Assume that the functions $\varphi_i : R_+ \to R_+$ have the form $\varphi_i(r) = 1 + r$ and the functions $a_i \in C(I)$. Denote by $b_i = \| a_i \| = \max[| a_i(t,s) | : t, s \in I]$. Then,

$$| f_i(t,s,x) | \leq a_i(t,s)(1 + | x |) \leq a_i(t,s) + b_i | x | .$$

Notice that this assumption is a special case of assumption (iii).

Consider now the assumptions $(iii)^*$ and $(vii)^*$ having the form
$(iii)^*$ $f_i : I \times R \to R$ are continuous and satisfy the Lipschitz condition

$$| f_i(t,s,x) - f_i(t,s,y) | \leq b_i | x - y |, \quad i = 1, 2.$$

From this assumption we can deduce that

$$| f_i(t,s,x) | - | f_i(t,s,0) | \leq | f_i(t,s,x) - f_i(t,s,0) | \leq b_i | x |$$

which implies that

$$| f_i(t,s,x) | \leq | f_i(t,s,0) | + b_i | x | = | a_i(t,s) | + b_i | x |$$

$(vii)^*$ $\mu b < 1$.

Corollary 2.4. Let the assumptions $(i) - (ii), (iii)^*, (iv) - (vi)$ and $(vii)^*$ be satisfied, then the coupled system (2) has an unique solution $(x, y) \in X$.

Proof. Let $u_1 = (x_1, y_1)$ and $u_2 = (x_2, y_2)$ be two solutions of the coupled system (2), we have

$$\begin{aligned} \| (x_1, y_1) - (x_2, y_2) \|_X &= \| (x_1 - x_2, y_1 - y_2) \|_X \\ &= \max\{\| x_1 - x_2 \|, \| y_1 - y_2 \|\} \end{aligned}$$

Now,

$$\begin{aligned} | x_1 - x_2 | &= | p_1(t) + \int_0^1 f_1(t,s,y_1(\psi_1(s))) \, d_s g_1(t,s) - p_1(t) + \int_0^1 f_1(t,s,y_2(\psi_1(s))) \, d_s g_1(t,s) | \\ &\leq \int_0^1 | f_1(t,s,y_1(\psi_1(s))) - f_1(t,s,y_2(\psi_1(s))) | \; d_s(\bigvee_{z=0}^{s} g_1(t,z)) \\ &\leq \int_0^1 b_1 | y_1(\psi_1(s)) - y_2(\psi_1(s)) | \; d_s g_1(t,s) \\ &\leq b \| y_1 - y_2 \| \int_0^1 d_s g_1(t,s) \\ &\leq b \| y_1 - y_2 \| [g_1(t,1) - g_1(t,0)] \\ &\leq b \| y_1 - y_2 \| [| g_1(t,1) | + | g_1(t,0) |] \\ &\leq \mu b \| y_1 - y_2 \| \end{aligned}$$

Unauthenticated
Download Date | 3/3/18 2:38 PM

Therefore,

$$\| x_1 - x_2 \| \leq \mu b \| y_1 - y_2 \| .$$

Also

$$\begin{aligned}
| y_1 - y_2 | &= | p_2(t) + \int_0^1 f_2(s, x_1(\psi_2(s))) \, d_s g_2(t,s) - p_2(t) + \int_0^1 f_2(s, x_2(\psi_2(s))) \, d_s g_2(t,s) | \\
&\leq \int_0^1 | f_2(s, x_1(\psi_2(s))) - f_2(s, x_2(\psi_2(s))) | \; d_s(\bigvee_{z=0}^{s} g_2(t,z)) \\
&\leq \int_0^1 b_2 | x_1(\psi_2(s)) - x_2(\psi_2(s)) | \; d_s g_2(t,s) \\
&\leq b \| x_1 - x_2 \| \int_0^t d_s g_2(t,s) \\
&\leq b \| x_1 - x_2 \| [g_2(t,1) - g_2(t,0)] \\
&\leq b \| x_1 - x_2 \| [| g_2(t,1) | + | g_2(t,0) |] \\
&\leq \mu b \| x_1 - x_2 \| .
\end{aligned}$$

Hence

$$\| y_1 - y_2 \| \leq \mu b \| x_1 - x_2 \| .$$

Then

$$\begin{aligned}
\| (x_1, y_1) - (x_2, y_2) \|_X &= \max\{\| x_1 - x_2 \|, \| y_1 - y_2 \|\} \\
&\leq \max\{\mu b \| y_1 - y_2 \|, \mu b \| x_1 - x_2 \|\} \\
&\leq \mu b \max\{\| y_1 - y_2 \|, \| x_1 - x_2 \|\} \\
&= \mu b \| (x_1, y_1) - (x_2, y_2) \|_X
\end{aligned}$$

which implies that

$$(1 - \mu b) \| (x_1, y_1) - (x_2, y_2) \|_X \leq 0,$$

therefore,

$$\| (x_1, y_1) - (x_2, y_2) \|_X = 0$$

This means that

$$(x_1, y_1) = (x_2, y_2) \quad \Rightarrow \quad x_1 = x_2, \quad y_1 = y_2.$$

Thus, the solution of the coupled system (2) is unique.

Example 3. Similarly as above, take the functions $a_i(t,s) \in C(I)$. Let us take the functions $\varphi_i : R_+ \to R_+$ having the form $\varphi_i(r) = 1 + r^\alpha$, where $\alpha > 0$ is a fixed number. Then

$$| f_i(t,s,x) | \leq a_i(t,s)(1 + | x |^\alpha) \leq a_i(t,s) + b_i | x |^\alpha,$$

where $b_i = \| a_i \|$.

Unauthenticated
Download Date | 3/3/18 2:38 PM

References

[1] J. Banaś, Some properties of Urysohn-Stieltjes integral operators, *Intern. J. Math. and Math. Sci.* 21(1998) 78-88.

[2] J. Banaś and J. Dronka, Integral operators of Volterra-Stieltjes type, their properties and applications, *Math. Comput. Modelling.* 32(2000) 1321-1331.

[3] J. Banaś, J.C. Mena, Some properties of nonlinear Volterra-Stieltjes integral operators, *Comput Math. Appl.* 49(2005) 1565-1573.

[4] J. Banaś, D. O'Regan, Volterra-Stieltjes integral operators, *Math. Comput. Modelling.* 41(2005) 335-344.

[5] J. Banaś, J.R. Rodriguez and K. Sadarangani, On a class of Urysohn-Stieltjes quadratic integral equations and their applications, *J. Comput. Appl. Math.* I13(2000) 35-50.

[6] J. Banaś and K. Sadarangani, Solvability of Volterra-Stieltjes operator-integral equations and their applications, *Comput Math. Appl.* 41(12)(2001) 1535-1544.

[7] C.W. Bitzer, Stieltjes-Volterra integral equations, *Illinois J. Math.* 14(1970) 434-451.

[8] S. Chen, Q. Huang and L.H. Erbe, Bounded and zero-convergent solutions of a class of Stieltjes integro-differential equations, *Proc. Amer. Math. Soc.* 113(1991) 999-1008.

[9] R.F. Curtain, A.J. Pritchard,: Functional analysis in modern applied mathematics. Academic press, London (1977).

[10] A.M.A. El-Sayed, H.H.G. Hashem, Existence results for coupled systems of quadratic integral equations of fractional orders, *Optimization Letters*, 7(2013) 1251-1260.

[11] A.M.A. El-Sayed, H.H.G. Hashem, Solvability of coupled systems of fractional order integro-differential equations, *J. Indones. Math. Soc.* 19(2)(2013) 111-121.

[12] H.H.G. Hashem, On successive approximation method for coupled systems of Chandrasekhar quadratic integral equations, *Journal of the Egyptian Mathematical Society.* 23(2015) 108-112.

[13] J.S. Macnerney, Integral equations and semigroups, *Illinois J. Math.* 7(1963) 148-173.

[14] A.B. Mingarelli, Volterra-Stieltjes integral equations and generalized ordinary differential expressions, *Lecture Notes in Math.*, 989, Springer (1983).

[15] I.P. Natanson, Theory of functions of a real variable, *Ungar, New York.* (1960).

Unauthenticated
Download Date | 3/3/18 2:38 PM